

ORIGINAL

Application Based on

Docket **83590AEK**

Inventors: Donald R. Diehl, Robert J. Nigier, Hans F. Schmitthenner,  
William J. Sonnefeld and Richard P. Dunlap  
Customer No. 01333

## SYNTHESIS OF PEPTOID SUBSTITUTED AZOLE COMPOUNDS

Commissioner for Patents,  
ATTN: BOX PATENT APPLICATION  
Washington, D. C. 20231

Express Mail Label No.: EL809162546US

Date: December 12, 2001

## SYNTHESIS OF PEPTOID SUBSTITUTED AZOLE COMPOUNDS

### **CROSS-REFERENCE TO RELATED APPLICATIONS**

- This application is related an application cofiled herewith under  
5 Attorney Docket No. 83554/AEK and directed to a photographic element, coupler  
and dye.

### **FIELD OF THE INVENTION**

- This invention relates to the synthesis of peptoid ballasted dye-  
10 forming azole couplers using a solid phase organic chemistry method of synthesis  
including reacting an amino functionalized azole compound with a resin bound  
peptoid oligomer bearing a terminal halogen substituent followed by cleavage of  
the resultant product from the resin surface using a fluorinated organic acid in an  
inert solvent.

15

### **BACKGROUND OF THE INVENTION**


- In the photographic art color images are formed by the exposure of  
a silver halide light sensitive element containing organic dye forming couplers  
followed by processing in aqueous developer solutions containing an aromatic  
20 primary amine color developing agent. The resultant dyes formed are yellow,  
magenta, or cyan and result in the formation of a color image in the photographic  
element.

- In the ink jet art images are formed by the deposition of a  
preformed colored ink or pigment upon a receiver material by use of a cartridge  
25 head loaded with a solution or dispersion of preformed dyes or pigments. The  
resultant images are colored as a result of the deposition of a varied mixture of the  
preformed inks or pigments.


- Both of these imaging methods require novel color forming  
materials to improve the properties of the resultant images. Color forming image  
30 dyes must have desirable hues and good stability with little or no fading or  
discoloration under storage in the dark or when exposed to light. The couplers  
which form these dyes should have good coupling efficiency and lead to dye

images with high contrast and high density in areas of maximum exposure and low density in areas of minimum exposure.

Couplers which have been recently received considerable interest in the imaging art include the pyrazolo[5,1-c]-1,2,4-triazole and the pyrazolo[1,5-b]-1,2,4-triazole heterocycles as shown below.




Pyrazolo[5,1-c]-1,2,4-triazole




Pyrazolo[1,5-b]-1,2,4-triazole



Other couplers include the imidazole, benzimidazole, and pyrroloazole couplers such as:



Imidazo[1,2-b]pyrazole



Pyrazolo[1,5-a]benzimidazole




Pyrrolo[1,2-b][1,2,4]triazoles

Other examples are the pyrazolone couplers as discussed hereinafter.


The R groups are substituents and X is H or a coupling-off group. It is well recognized by those skilled in the art that modifications of the structure of the substituents on the pyrazolotriazole heterocycle can have important effects on the dyes formed from these couplers, including improvements in dye image stability, etc. Thus, the researcher skilled in the art of photographic couplers is always seeking new substituents on the pyrazolotriazole heterocycle with an aim to discover ever better image dye properties. In recent years many patents have been issued on variations of the 3-alkyl substituted pyrazolo[5,1-c]-1,2,4-triazole couplers. These couplers have the required hue, contrast, and efficiency properties necessary to produce superior image dyes in modern photographic materials. Nevertheless, new substituent research continues.

In Bailey's original patent, US 3,810,761 (5/1974), a wide range of substituents are described as useful in varying the hue and properties of pyrazolo[5,1-c]-1,2,4-triazole image dyes. In Harder's patent, US 5,925,503 (7/1999), a tetramethylethyl link in the 3-position of the pyrazolo[5,1-c]-1,2,4-triazole is reported to provide image dyes with superior light stability and hue. In Harder's patented structure, see Coupler Structure A below, W is reported to represent  $C(O)R^5$ ,  $SO_2R^5$ , or  $P(O)(OR^6)_2$ .

The properties of a pyrazolo[5,1-c]-1,2,4-triazole coupler similar to Structure A in which W is reported to be a CHR structural unit would be of interest to the skilled photographic coupler researcher. However, a synthesis method for such a pyrazolo[5,1-c]-1,2,4-triazole coupler is unknown. This is because the alkylation of the pendant amino functional group in Structure A by typical alkylating agents such as methyl iodide, ethyl iodide, methyltosylate, etc., results in the simultaneous alkylation of the ring nitrogen centers at position 1 and position 5 of the pyrazolo[5,1-c]-1,2,4-triazole ring thus producing products which contain multiple alkylated centers. Thus, the synthesis of new and improved pyrazolo[5,1-c]-1,2,4-triazoles has been limited in the past to those materials of the type described in Harder US 5,925,503 (7/1999).



10



83590US1/AEK

It has now been found that the use of the alkylation protocol as reported above does not result in the multiple alkylation of the ring nitrogen centers at position 1 and position 5 of the pyrazoloazole ring as demonstrated by such typical alkylating agents as methyl iodide, ethyl iodide, and methyltosylate, etc., but instead, results in the alkylation of the pendant amino functional group shown in Coupler Structure A without further alkylation occurring. This straightforward synthesis enables the production for the first time of pyrazoloazole couplers bearing peptoid substituents in the W position of Coupler Structure A and enables the production of photographic elements and dyes using the couplers.


It would be useful to have photographic materials containing novel coupler compounds that produce dyes that exhibit desirable spectral absorption characteristics and good stability.

#### SUMMARY OF THE INVENTION


The invention provides a process for the preparation of a peptoid substituted azole compound comprising reacting an amino functionalized azole compound with a resin bound peptoid oligomer bearing a terminal halogen substituent followed by cleavage of the resultant product from the resin surface using a fluorinated organic acid in an inert solvent

#### DETAILED DESCRIPTION OF THE INVENTION

A detailed description of the invention is provided as follows. A peptoid group is a functionality depicted below as defined by R.N. Zuckermann, et. al., *J. Am. Chem. Soc.*, **1992**, 10646-10647 in which R is generally defined as a side chain and X as an  $\text{NH}_2$ , and n is a number equal to or greater than zero,




or, for purposes of this invention,




5 The couplers useful in the element of the invention contain a peptoid group represented by the above formula. The couplers form dyes having desirable hue and stability for imaging. They conveniently contain up to 7, suitably 2-4 acetamide groups, and they are desirably linked to the coupler nucleus by a tetramethyl ethyl group.


10 Azole couplers are well-known and include one ring as in the pyrazolones or two fused 5-membered rings at least one being an azole. Azole couplers include, for example, the pyrazolo[5,1-c]-1,2,4-triazole, pyrazolo[1,5-b]-1,2,4-triazole, imidazo[1,2-b]pyrazole, pyrazolo[1,5-a]benzimidazole and pyrrolo[1,2-b][1,2,4] triazole heterocycles as shown below in which each R group is a  
15 substituent and X is H or a coupling off group.




Pyrazolo[5,1-c]-1,2,4-triazole





Pyrazolo[1,5-b]-1,2,4-triazole



Imidazo[1,2-b]pyrazole




Pyrazolo[1,5-a]benzimidazole



# Pyrrolo[1,2-b][1,2,4]triazoles

5 The R groups are substituents and X is H or a coupling-off group. More commonly, the second ring is a triazole such as the 1H-pyrazolo[5,1-c]-1,2,4-triazole and a 1H-pyrazolo[1,5-b]-1,2,4-triazole, as shown above.

A generic formula that represents a suitably substituted bicyclic azole coupler is shown in Formula I below:



I

wherein

each R1 represents an independently selected substituent group and a is 0-4;

each R2 represents an independently selected substituent group;

each R3 represents hydrogen or an independently selected substituent group;

L represents a single bond or a chain of atoms containing one or more of carbon, nitrogen, oxygen, and sulfur atoms;

each Y, Y', and Y'' independently represents a carbon or nitrogen atom with the proviso that at least one of Y, Y' or Y'' is a carbon, and that Y and Y' may represent the carbons necessary to form part of a fused six membered



aromatic ring, and provided further that, if Y and Y' represents the carbons necessary to form part of a fused six membered aromatic ring, the L linking group is not attached directly to the Y or Y' atom, and Y, Y' and Y'' may be further substituted by R1 when not directly attached to L;

X is a hydrogen atom, a halogen atom, a carboxy group, an acyl group, or a group bonded to the coupling position through an oxygen, nitrogen, or sulfur atom, and

Z is 1-6.

Desirably a is at least 1 and each R1 represents a substituent such as those listed hereinafter. Typical examples of such groups include methyl, ethyl, butyl, dodecyl, octadecyl, icosenyl, iso-propyl, tert-butyl, tert-octyl, tert-dodecyl, cyclohexyl, cyclopentyl, allyl, vinyl, 2-hexadecenyl, and propargyl groups. Especially preferred R1 substituents are alkyl, aryl, alkoxy, amino, anilino, alkoxycarbonyl, carbamoyl, acyl, cyano, sulfone, and sulfonamido groups and particularly t-butyl, phenyl, ethylcarboxyl, methylcyclopropyl, and adamantyl groups.

R2 may be a substituent such as an alkyl, cycloalkyl, aryl, or heteroaryl group. Useful examples include groups containing up to 20 carbon atoms such as cyclohexyl, 2-ethylhexyl, dodecyl, 3-ethoxypropyl, cyclohexylmethyl, benzyl, phenyl, and 1-(3-propyl)-2-pyrrolidinone groups.

R3 may be hydrogen or a substituent group such as one of the substituent groups specified for R1. Hydrogen is conveniently employed


The group X is useful as hydrogen or a leaving (or coupling-off) group. Examples of groups represented by X include but are not limited to hydrogen or any of the coupling off groups known in the art as described more fully hereinafter. Especially preferred X groups include hydrogen and halogen.

The atoms represented by Y, Y', and Y'' are nitrogen or carbon. At least one of them is carbon and Y and Y' may join to form a fused six-membered aromatic ring so long as the group L is not bonded to Y or Y'.

The value of "Z" is indicative of the length of the peptoid chain. Typically Z is 1-3.

Azomethine dyes of the invention are formed upon reaction of an oxidized primary aromatic amine with the azole couplers useful in this invention. Preferred dyes of this invention upon suitable dispersion preparation and ink jet coating on a suitable receiver provide magenta images with a spectral absorption peak in the region of the spectrum from 500 to 600 nm with a half band width of less then 120 nm, preferably less than 100 nm and most preferably less than 90 nm and exhibit the property of good light and dark stability. The couplers of this invention may also optionally be prepared into suitable dispersions which when coated upon a support bearing a light sensitive silver halide emulsion and further after exposure and processing in an oxidized primary aromatic amine developing solution may produce magenta images with a spectral absorption peak in the region of the spectrum from 500 to 600 nm with a half band width of less then 120 nm, preferably less than 100 nm and most preferably less than 90 nm and exhibit the property of good light and dark stability.


The following are examples of couplers useful in the invention:



A

Invention Coupler	R <sub>1</sub>	R <sub>2</sub>	X
A1	t-Bu	cyclohexyl	H
A2	t-Bu	2-ethylhexyl	H
A3	t-Bu	dodecyl	H
A4	t-Bu	3-ethoxypropyl	H
A5	t-Bu	cyclohexanemethyl	H
A6	t-Bu	benzyl	H


Invention Coupler	R <sub>1</sub>	R <sub>2</sub>	X
A7	t-Bu	phenyl	H
A8	t-Bu	1-(3-propyl) -2-pyrrolidinone	H
A9	t-Bu	cyclohexyl	Cl
A10	t-Bu	2-ethylhexyl	Cl
A11	t-Bu	dodecyl	Cl
A12	t-Bu	3-ethoxypropyl	Cl
A13	t-Bu	cyclohexanemethyl	Cl
A14	t-Bu	benzyl	Cl
A15	Ph	cyclohexyl	H
A16	Ph	2-ethylhexyl	H
A17	Ph	dodecyl	H
A18	Ph	3-ethoxypropyl	H
A19	Ph	cyclohexanemethyl	H
A20	Ph	benzyl	H
A21	CO <sub>2</sub> Et	cyclohexyl	H
A22	CO <sub>2</sub> Et	2-ethylhexyl	H
A23	CO <sub>2</sub> Et	dodecyl	H
A24	CO <sub>2</sub> Et	3-ethoxypropyl	H
A25	CO <sub>2</sub> Et	cyclohexanemethyl	H
A26	CO <sub>2</sub> Et	benzyl	H



A'

5


Invention Coupler	R <sub>1</sub>	R <sub>2</sub>	X
A'1	t-Bu	cyclohexyl	H
A'2	t-Bu	2-ethylhexyl	H
A'3	t-Bu	dodecyl	H
A'4	t-Bu	3-ethoxypropyl	H
A'5	t-Bu	cyclohexanemethyl	H
A'6	t-Bu	benzyl	H
A'7	t-Bu	phenyl	H
A'8	t-Bu	1-(3-propyl) -2-pyrrolidinone	H
A'9	t-Bu	cyclohexyl	Cl
A'10	t-Bu	2-ethylhexyl	Cl
A'11	t-Bu	dodecyl	Cl
A'12	t-Bu	3-ethoxypropyl	Cl
A'13	t-Bu	cyclohexanemethyl	Cl
A'14	t-Bu	benzyl	Cl
A'15	Me	cyclohexyl	H
A'16	Me	2-ethylhexyl	H
A'17	Me	dodecyl	H
A'18	Me	3-ethoxypropyl	H
A'19	Me	cyclohexanemethyl	H
A'20	Me	benzyl	H



B

Invention Coupler	$R_1$	$R_2$	$R_2'$	X
B1	t-Bu	cyclohexyl	cyclohexyl	H
B2	t-Bu	2-ethylhexyl	cyclohexyl	H
B3	t-Bu	dodecyl	cyclohexyl	H
B4	t-Bu	cyclohexyl	2-ethylhexyl	H
B5	t-Bu	2-ethylhexyl	2-ethylhexyl	H
B6	t-Bu	dodecyl	2-ethylhexyl	H
B7	t-Bu	cyclohexyl	dodecyl	H
B8	t-Bu	2-ethylhexyl	dodecyl	H
B9	t-Bu	dodecyl	dodecyl	H
B10	t-Bu	hexyl	hexyl	H
B11	t-Bu	3-ethoxypropyl	hexyl	H
B12	t-Bu	cyclohexanemethyl	hexyl	H
B13	t-Bu	benzyl	hexyl	H
B14	t-Bu	hexyl	3-ethoxypropyl	H
B15	t-Bu	3-ethoxypropyl	3-ethoxypropyl	H
B16	t-Bu	cyclohexanemethyl	3-ethoxypropyl	H
B17	t-Bu	benzyl	3-ethoxypropyl	H
B18	t-Bu	hexyl	cyclohexanemethyl	H
B19	t-Bu	3-ethoxypropyl	cyclohexanemethyl	H
B20	t-Bu	cyclohexanemethyl	cyclohexanemethyl	H
B21	t-Bu	benzyl	cyclohexanemethyl	H
B22	t-Bu	hexyl	benzyl	H
B23	t-Bu	3-ethoxypropyl	benzyl	H
B24	t-Bu	cyclohexanemethyl	benzyl	H

Invention Coupler	R <sub>1</sub>	R <sub>2</sub>	R <sub>2</sub> '	X
B25	t-Bu	benzyl	benzyl	H
B26	t-Bu	cyclohexyl	cyclohexyl	Cl
B27	t-Bu	2-ethylhexyl	cyclohexyl	Cl
B28	t-Bu	dodecyl	cyclohexyl	Cl
B29	Ph	cyclohexyl	cyclohexyl	H
B30	Ph	2-ethylhexyl	cyclohexyl	H
B31	Ph	dodecyl	cyclohexyl	H
B32	CO <sub>2</sub> Et	cyclohexyl	cyclohexyl	H
B33	CO <sub>2</sub> Et	2-ethylhexyl	cyclohexyl	H
B34	CO <sub>2</sub> Et	dodecyl	cyclohexyl	H




**B'**

5

Invention Coupler	R <sub>1</sub>	R <sub>2</sub>	R <sub>2</sub> '	X
B'1	t-Bu	cyclohexyl	cyclohexyl	H
B'2	t-Bu	2-ethylhexyl	cyclohexyl	H
B'3	t-Bu	dodecyl	cyclohexyl	H
B'4	t-Bu	cyclohexyl	2-ethylhexyl	H
B'5	t-Bu	2-ethylhexyl	2-ethylhexyl	H
B'6	t-Bu	dodecyl	2-ethylhexyl	H
B'7	t-Bu	cyclohexyl	dodecyl	H
B'8	t-Bu	2-ethylhexyl	dodecyl	H
B'9	t-Bu	dodecyl	dodecyl	H

Invention Coupler	R <sub>1</sub>	R <sub>2</sub>	R <sub>2</sub> '	X
B'10	t-Bu	hexyl	hexyl	H
B'11	t-Bu	3-ethoxypropyl	hexyl	H
B'12	t-Bu	cyclohexanemethyl	hexyl	H
B'13	t-Bu	benzyl	hexyl	H
B'14	t-Bu	hexyl	3-ethoxypropyl	H
B'15	t-Bu	3-ethoxypropyl	3-ethoxypropyl	H
B'16	t-Bu	cyclohexanemethyl	3-ethoxypropyl	H
B'17	t-Bu	benzyl	3-ethoxypropyl	H
B'18	t-Bu	hexyl	cyclohexanemethyl	H
B'19	t-Bu	3-ethoxypropyl	cyclohexanemethyl	H
B'20	t-Bu	cyclohexanemethyl	cyclohexanemethyl	H
B'21	t-Bu	benzyl	cyclohexanemethyl	H
B'22	t-Bu	hexyl	benzyl	H
B'23	t-Bu	3-ethoxypropyl	benzyl	H
B'24	t-Bu	cyclohexanemethyl	benzyl	H
B'25	t-Bu	benzyl	benzyl	H
B'26	t-Bu	cyclohexyl	cyclohexyl	Cl
B'27	t-Bu	2-ethylhexyl	cyclohexyl	Cl
B'28	t-Bu	dodecyl	cyclohexyl	Cl
B'29	Me	cyclohexyl	cyclohexyl	H
B'30	Me	2-ethylhexyl	cyclohexyl	H
B'31	Me	dodecyl	cyclohexyl	H




C

Invention Coupler	R <sub>1</sub>	R <sub>2</sub>	R <sub>2</sub> '	R <sub>2</sub> ''	X
C1	t-Bu	cyclohexyl	cyclohexyl	cyclohexyl	H
C2	t-Bu	2-ethylhexyl	cyclohexyl	2-ethylhexyl	H
C3	t-Bu	dodecyl	cyclohexyl	dodecyl	H
C4	t-Bu	cyclohexyl	2-ethylhexyl	cyclohexyl	H
C5	t-Bu	2-ethylhexyl	2-ethylhexyl	2-ethylhexyl	H
C6	t-Bu	dodecyl	2-ethylhexyl	dodecyl	H
C7	t-Bu	cyclohexyl	dodecyl	cyclohexyl	H
C8	t-Bu	2-ethylhexyl	dodecyl	2-ethylhexyl	H
C9	t-Bu	dodecyl	dodecyl	dodecyl	H
C10	t-Bu	hexyl	hexyl	hexyl	H
C11	t-Bu	3-ethoxypropyl	hexyl	3-ethoxypropyl	H
C12	t-Bu	cyclohexanemethyl	hexyl	cyclohexanemethyl	H
C13	t-Bu	benzyl	hexyl	benzyl	H
C14	t-Bu	hexyl	3-ethoxypropyl	hexyl	H
C15	t-Bu	3-ethoxypropyl	3-ethoxypropyl	3-ethoxypropyl	H
C16	t-Bu	cyclohexanemethyl	3-ethoxypropyl	cyclohexanemethyl	H
C17	t-Bu	benzyl	3-ethoxypropyl	benzyl	H
C18	t-Bu	hexyl	cyclohexanemethyl	hexyl	H
C19	t-Bu	3-ethoxypropyl	cyclohexanemethyl	3-ethoxypropyl	H
C20	t-Bu	cyclohexanemethyl	cyclohexanemethyl	cyclohexanemethyl	H
C21	t-Bu	benzyl	cyclohexanemethyl	benzyl	H
C22	t-Bu	hexyl	benzyl	hexyl	H
C23	t-Bu	3-ethoxypropyl	benzyl	3-ethoxypropyl	H



Invention Coupler	R <sub>1</sub>	R <sub>2</sub>	R <sub>2</sub> '	R <sub>2</sub> ''	X
C24	t-Bu	cyclohexanemethyl	benzyl	cyclohexanemethyl	H
C25	t-Bu	benzyl	benzyl	benzyl	H
C26	t-Bu	cyclohexyl	cyclohexyl	cyclohexyl	Cl
C27	t-Bu	2-ethylhexyl	cyclohexyl	2-ethylhexyl	Cl
C28	t-Bu	dodecyl	cyclohexyl	dodecyl	Cl
C29	Ph	cyclohexyl	cyclohexyl	cyclohexyl	H
C30	Ph	2-ethylhexyl	cyclohexyl	2-ethylhexyl	H
C31	Ph	dodecyl	cyclohexyl	dodecyl	H
C32	CO <sub>2</sub> Et	cyclohexyl	cyclohexyl	cyclohexyl	H
C33	CO <sub>2</sub> Et	2-ethylhexyl	cyclohexyl	2-ethylhexyl	H
C34	CO <sub>2</sub> Et	dodecyl	cyclohexyl	dodecyl	H




5

## D


Invention Coupler	R <sub>1</sub>	R <sub>2</sub>	R <sub>2</sub> '	X
D1	t-Bu	hexyl	hexyl	H
D2	t-Bu	3-ethoxypropyl	hexyl	H
D3	t-Bu	cyclohexanemethyl	hexyl	H
D4	t-Bu	benzyl	hexyl	H
D5	t-Bu	hexyl	3-ethoxypropyl	H
D6	t-Bu	3-ethoxypropyl	3-ethoxypropyl	H
D7	t-Bu	cyclohexanemethyl	3-ethoxypropyl	H
D8	t-Bu	benzyl	3-ethoxypropyl	H

Invention Coupler	R <sub>1</sub>	R <sub>2</sub>	R <sub>2</sub> '	X
D9	t-Bu	hexyl	cyclohexanemethyl	H
D10	t-Bu	3-ethoxypropyl	cyclohexanemethyl	H
D11	t-Bu	cyclohexanemethyl	cyclohexanemethyl	H
D12	t-Bu	benzyl	cyclohexanemethyl	H
D13	t-Bu	hexyl	benzyl	H
D14	t-Bu	3-ethoxypropyl	benzyl	H
D15	t-Bu	cyclohexanemethyl	benzyl	H
D16	t-Bu	benzyl	benzyl	H
D17	t-Bu	cyclohexyl	cyclohexyl	H
D18	t-Bu	2-ethylhexyl	cyclohexyl	H
D19	t-Bu	dodecyl	cyclohexyl	H
D20	t-Bu	cyclohexyl	2-ethylhexyl	H
D21	t-Bu	2-ethylhexyl	2-ethylhexyl	H
D22	t-Bu	dodecyl	2-ethylhexyl	H
D23	t-Bu	cyclohexyl	dodecyl	Cl
D24	t-Bu	2-ethylhexyl	dodecyl	Cl
D25	t-Bu	dodecyl	dodecyl	Cl
D26	t-Bu	dodecyl	hexyl	Cl
D27	t-Bu	dodecyl	3-ethoxypropyl	H
D28	t-Bu	dodecyl	cyclohexanemethyl	H
D29	t-Bu	2-ethylhexyl	hexyl	H
D30	t-Bu	2-ethylhexyl	3-ethoxypropyl	H
D31	t-Bu	2-ethylhexyl	cyclohexanemethyl	H
D32	t-Bu	cyclohexyl	hexyl	H
D33	t-Bu	cyclohexyl	3-ethoxypropyl	H
D34	t-Bu	cyclohexyl	cyclohexanemethyl	H



E


Invention Coupler	R <sub>1</sub>	R <sub>2</sub>	X
E1	t-Bu	cyclohexyl	H
E2	t-Bu	2-ethylhexyl	H
E3	t-Bu	dodecyl	H
E4	t-Bu	3-ethoxypropyl	H
E5	t-Bu	cyclohexanemethyl	H
E6	t-Bu	benzyl	H
E7	t-Bu	phenyl	H
E8	t-Bu	1-(3-propyl) -2-pyrrolidinone	H
E9	t-Bu	cyclohexyl	Cl
E10	t-Bu	2-ethylhexyl	Cl
E11	t-Bu	dodecyl	Cl
E12	t-Bu	3-ethoxypropyl	Cl
E13	t-Bu	cyclohexanemethyl	Cl
E14	t-Bu	benzyl	Cl



F

Invention Coupler	R <sub>1</sub>	R <sub>2</sub>	R <sub>2</sub> '	X
F1	t-Bu	hexyl	hexyl	H
F2	t-Bu	3-ethoxypropyl	hexyl	H
F3	t-Bu	cyclohexanemethyl	hexyl	H
F4	t-Bu	benzyl	hexyl	H
F5	t-Bu	hexyl	3-ethoxypropyl	H
F6	t-Bu	3-ethoxypropyl	3-ethoxypropyl	H
F7	t-Bu	cyclohexanemethyl	3-ethoxypropyl	H
F8	t-Bu	benzyl	3-ethoxypropyl	H
F9	t-Bu	hexyl	cyclohexanemethyl	H
F10	t-Bu	3-ethoxypropyl	cyclohexanemethyl	H
F11	t-Bu	cyclohexanemethyl	cyclohexanemethyl	H
F12	t-Bu	benzyl	cyclohexanemethyl	H
F13	t-Bu	hexyl	benzyl	H
F14	t-Bu	3-ethoxypropyl	benzyl	H
F15	t-Bu	cyclohexanemethyl	benzyl	H
F16	t-Bu	benzyl	benzyl	H
F17	t-Bu	cyclohexyl	cyclohexyl	H
F18	t-Bu	2-ethylhexyl	cyclohexyl	H
F19	t-Bu	dodecyl	cyclohexyl	H
F20	t-Bu	cyclohexyl	2-ethylhexyl	H
F21	t-Bu	2-ethylhexyl	2-ethylhexyl	H
F22	t-Bu	dodecyl	2-ethylhexyl	H
F23	t-Bu	cyclohexyl	dodecyl	Cl

Invention Coupler	R <sub>1</sub>	R <sub>2</sub>	R <sub>2</sub> '	X
F24	t-Bu	2-ethylhexyl	dodecyl	Cl
F25	t-Bu	dodecyl	dodecyl	Cl
F26	t-Bu	dodecyl	hexyl	Cl
F27	t-Bu	dodecyl	3-ethoxypropyl	H
F28	t-Bu	dodecyl	cyclohexanemethyl	H
F29	t-Bu	2-ethylhexyl	hexyl	H
F30	t-Bu	2-ethylhexyl	3-ethoxypropyl	H
F31	t-Bu	2-ethylhexyl	cyclohexanemethyl	H
F32	t-Bu	cyclohexyl	hexyl	H
F33	t-Bu	cyclohexyl	3-ethoxypropyl	H
F34	t-Bu	cyclohexyl	cyclohexanemethyl	H




PT

Invention Coupler	R <sub>1</sub>	R <sub>1</sub> '	R <sub>2</sub>	R <sub>2</sub> '	X
PT1	CN	CO <sub>2</sub> Et	hexyl	hexyl	H
PT2	CN	CO <sub>2</sub> Et	3-ethoxypropyl	hexyl	H
PT3	CN	CO <sub>2</sub> Et	cyclohexanemethyl	hexyl	H
PT4	CN	CO <sub>2</sub> Et	benzyl	hexyl	H
PT5	CN	CO <sub>2</sub> Et	hexyl	3-ethoxypropyl	H
PT6	CN	CO <sub>2</sub> Et	3-ethoxypropyl	3-ethoxypropyl	H
PT7	CN	CO <sub>2</sub> Et	cyclohexanemethyl	3-ethoxypropyl	H
PT8	CN	CO <sub>2</sub> Et	benzyl	3-ethoxypropyl	H
PT9	CN	CN	hexyl	cyclohexanemethyl	H

Invention Coupler	R <sub>1</sub>	R <sub>1</sub> '	R <sub>2</sub>	R <sub>2</sub> '	X
PT10	CN	CN	3-ethoxypropyl	cyclohexanemethyl	H
PT11	CN	CN	cyclohexanemethyl	cyclohexanemethyl	H
PT12	CN	CN	benzyl	cyclohexanemethyl	H
PT13	CN	CN	hexyl	benzyl	H
PT14	CN	CN	3-ethoxypropyl	benzyl	H
PT15	CN	CN	cyclohexanemethyl	benzyl	H
PT16	CN	CN	benzyl	benzyl	H
PT17	CONHPh	CN	2-ethylhexyl	cyclohexanemethyl	H
PT18	CONHPh	CN	dodecyl	cyclohexanemethyl	H
PT19	CONHPh	CN	cyclohexanemethyl	2-ethylhexyl	H
PT20	CONHPh	CN	2-ethylhexyl	2-ethylhexyl	H


The couplers of the present invention may be used as intermediates to prepare novel dyes. As described by Bailey in EP 1,253,933 such novel dyes may be useful in sensitized silver halide photographic materials as filter dyes or sensitizing dyes. Such dyes may also be useful as textile dyes and in more modern imaging applications such as a thermal dye transfer imaging processes and as ink jet dyes. For example, azomethine dyes of the generic Dye Structure GI are shown below.



Generic Dye Structure GI

Most especially preferred dyes are the azomethine dyes of generic Dye Structure II below. These dyes are easily prepared by the reaction of oxidized primary aromatic amines with the novel pyrazolo[5,1-c]-1,2,4-triazole couplers of this invention. These new dyes have good extinction and high optical

density with low unwanted absorbance at wavelengths of light which reduce the desirable hue of the dye. The azomethine dyes prepared from the couplers of this invention have been found to have good light stability. This particular stability may result from the intramolecular hydrogen bonding of the N1 heterocycle center of the pyrazolo[5,1-c]-1,2,4-triazole coupler with the terminal primary amide functionality of the peptoid substituent.



Generic Dye Structure II

One skilled in the art of image coupler synthesis may recognize that L groups in coupler Structure I may represent alkyl, aryl, or heteroaryl groups which may in turn be further substituted. Examples of such substituents may typically include those described below but are not limited by those listed.

The variables in the dye structures are defined as shown above for formula I.


The novel dyes of Dye Structure II may be represented by the couplers bearing a peptoid functionality as part of the 3-substituent on the pyrazolo[5,1-c]-1,2,4-triazole heterocycle.

In the novel Dye Structure II the substituents Rn represent one or more substituents defining typical aminophenol and p-phenylenediamine derivatives which may include but are not limited by the following: o-aminophenol, p-aminophenol, 5-amino-2-hydroxytoluene, 2-amino-3-hydroxytoluene, 2-hydroxy-3-amino-1,4-dimethylbenzene, N,N-diethyl-p-phenylenediamine, 2-amino-5-diethylaminotoluene, 2-amino-5-(N-ethyl-N-laurylamino)toluene, 4-[N-ethyl-N-(beta-hydroxyethyl)amino]aniline, 2-methyl-4-[N-ethyl-N-(beta-hydroxyethyl)amino]aniline, 4-amino-3-methyl-N-ethyl-N-

[beta-(methanesulfonamido)ethyl]aniline, N-(2-amino-5-diethylaminophenylethyl)methanesulfonamide, N,N-dimethyl-p-phenylenediamine monohydrochloride, 4-N,N-diethyl-2-methylphenylenediamine monohydrochloride, 4-(N-ethyl-N-2-methanesulfonylaminoethyl)-2-methylphenylenediamine sesquisulfate monohydrate, 4-(N-ethyl-N-2-hydroxyethyl)-2-methylphenylenediamine sulfate, 4-amino-3-methyl-N-ethyl-N-methoxyethylaniline, 4-amino-3-methyl-N-ethyl-N-beta-ethoxyethylaniline, 4-amino-3-methyl-N-ethyl-N-beta-butoxyethylaniline, and 4-N,N-diethyl-2,2'-methanesulfonylaminoethylphenylenediamine hydrochloride.

Particularly preferred are the p-phenylenediamines and especially the N,N-dialkyl-p-phenylenediamines in which the alkyl groups or the aromatic nucleus can be substituted or unsubstituted.

Hereunder, the typical examples of the azomethine dyes relating to the present invention will be given. However, the present invention shall not be limited thereto.




G


Invention Dye	Invention Coupler	R3	R4	R5
G1	A1	Me	Et	-CH <sub>2</sub> CH <sub>2</sub> NHSO <sub>2</sub> Me
G2	A2	Me	Et	-CH <sub>2</sub> CH <sub>2</sub> NHSO <sub>2</sub> Me
G3	A3	Me	Et	-CH <sub>2</sub> CH <sub>2</sub> NHSO <sub>2</sub> Me
G4	B1	Me	Et	-CH <sub>2</sub> CH <sub>2</sub> NHSO <sub>2</sub> Me
G5	B2	Me	Et	-CH <sub>2</sub> CH <sub>2</sub> NHSO <sub>2</sub> Me
G6	B3	Me	Et	-CH <sub>2</sub> CH <sub>2</sub> NHSO <sub>2</sub> Me
G7	B4	Me	Et	-CH <sub>2</sub> CH <sub>2</sub> NHSO <sub>2</sub> Me




<u>Invention</u> Dye	<u>Invention</u> Coupler	<u>R3</u>	<u>R4</u>	<u>R5</u>
G8	B5	Me	Et	-CH <sub>2</sub> CH <sub>2</sub> NHSO <sub>2</sub> Me
G9	B6	Me	Et	-CH <sub>2</sub> CH <sub>2</sub> NHSO <sub>2</sub> Me
G10	B7	Me	Et	-CH <sub>2</sub> CH <sub>2</sub> NHSO <sub>2</sub> Me
G11	B8	Me	Et	-CH <sub>2</sub> CH <sub>2</sub> NHSO <sub>2</sub> Me
G12	B9	Me	Et	-CH <sub>2</sub> CH <sub>2</sub> NHSO <sub>2</sub> Me
G13	B10	Me	Et	-CH <sub>2</sub> CH <sub>2</sub> NHSO <sub>2</sub> Me
G14	B11	Me	Et	-CH <sub>2</sub> CH <sub>2</sub> NHSO <sub>2</sub> Me
G15	B12	Me	Et	-CH <sub>2</sub> CH <sub>2</sub> NHSO <sub>2</sub> Me
G16	B13	Me	Et	-CH <sub>2</sub> CH <sub>2</sub> NHSO <sub>2</sub> Me
G17	B14	Me	Et	-CH <sub>2</sub> CH <sub>2</sub> NHSO <sub>2</sub> Me
G18	B15	Me	Et	-CH <sub>2</sub> CH <sub>2</sub> NHSO <sub>2</sub> Me
G19	B16	Me	Et	-CH <sub>2</sub> CH <sub>2</sub> NHSO <sub>2</sub> Me
G20	B17	Me	Et	-CH <sub>2</sub> CH <sub>2</sub> NHSO <sub>2</sub> Me
G21	B18	Me	Et	-CH <sub>2</sub> CH <sub>2</sub> NHSO <sub>2</sub> Me
G22	B19	Me	Et	-CH <sub>2</sub> CH <sub>2</sub> NHSO <sub>2</sub> Me
G23	B20	Me	Et	-CH <sub>2</sub> CH <sub>2</sub> NHSO <sub>2</sub> Me
G24	B21	Me	Et	-CH <sub>2</sub> CH <sub>2</sub> NHSO <sub>2</sub> Me
G25	B22	Me	Et	-CH <sub>2</sub> CH <sub>2</sub> NHSO <sub>2</sub> Me
G26	B23	Me	Et	-CH <sub>2</sub> CH <sub>2</sub> NHSO <sub>2</sub> Me
G27	B24	Me	Et	-CH <sub>2</sub> CH <sub>2</sub> NHSO <sub>2</sub> Me
G28	B25	Me	Et	-CH <sub>2</sub> CH <sub>2</sub> NHSO <sub>2</sub> Me
G29	D1	Me	Et	-CH <sub>2</sub> CH <sub>2</sub> NHSO <sub>2</sub> Me
G30	D2	Me	Et	-CH <sub>2</sub> CH <sub>2</sub> NHSO <sub>2</sub> Me
G31	D3	Me	Et	-CH <sub>2</sub> CH <sub>2</sub> NHSO <sub>2</sub> Me
G32	D4	Me	Et	-CH <sub>2</sub> CH <sub>2</sub> NHSO <sub>2</sub> Me
G33	D5	Me	Et	-CH <sub>2</sub> CH <sub>2</sub> NHSO <sub>2</sub> Me
G34	D6	Me	Et	-CH <sub>2</sub> CH <sub>2</sub> NHSO <sub>2</sub> Me
G35	D7	Me	Et	-CH <sub>2</sub> CH <sub>2</sub> NHSO <sub>2</sub> Me
G36	D8	Me	Et	-CH <sub>2</sub> CH <sub>2</sub> NHSO <sub>2</sub> Me




Invention Dye G37



Invention Dye G38



Invention Dye G39



Invention Dye G40

The couplers of this invention will be recognized by one skilled in the art as useful in photographic elements. The photographic elements of this invention can be single color elements or multicolor elements. Multicolor elements typically contain dye image-forming units sensitive to each of the three primary regions of the visible spectrum. Each unit can be comprised of a single emulsion layer or of multiple emulsion layers sensitive to a given region of the spectrum. The layers of the element, including the layers of the image-forming units, can be arranged in various orders as known in the art.

Unless otherwise specifically stated, use of the term "group", "substituted" or "substituent" means any group or atom other than hydrogen. Additionally, when the term "group" is used, it means that when a substituent group contains a substitutable hydrogen, it is also intended to encompass not only the substituent's unsubstituted form, but also its form further substituted with any substituent group or groups as herein mentioned, so long as the substituent does not destroy properties necessary for photographic utility. Suitably, a substituent group may be halogen or may be bonded to the remainder of the molecule by an atom of carbon, silicon, oxygen, nitrogen, phosphorous, or sulfur. The substituent may be, for example, halogen, such as chlorine, bromine or fluorine; nitro; hydroxyl; cyano; carboxyl; or groups which may be further substituted, such as alkyl, including straight or branched chain or cyclic alkyl, such as methyl, trifluoromethyl, ethyl, *t*-butyl, 3-(2,4-di-*t*-pentylphenoxy) propyl, cyclohexyl, and tetradecyl; alkenyl, such as ethylene, 2-butene; alkoxy, such as methoxy, ethoxy, propoxy, butoxy, 2-methoxyethoxy, *sec*-butoxy, hexyloxy, 2-ethylhexyloxy, tetradecyloxy, 2-(2,4-di-*t*-pentylphenoxy)ethoxy, and 2-dodecyloxyethoxy; aryl such as phenyl, 4-*t*-butylphenyl, 2,4,6-trimethylphenyl, naphthyl; aryloxy, such as phenoxy, 2-methylphenoxy, alpha- or beta-naphthylloxy, and 4-tolyloxy; carbonamido, such as acetamido, benzamido, butyramido, tetradecanamido, alpha-(2,4-di-*t*-pentyl-phenoxy)acetamido, alpha-(2,4-di-*t*-pentylphenoxy)butyramido, alpha-(3-pentadecylphenoxy)-hexanamido, alpha-(4-hydroxy-3-*t*-butylphenoxy)-tetradecanamido, 2-oxo-pyrrolidin-1-yl, 2-oxo-5-tetradecylpyrrolin-1-yl, N-methyltetradecanamido, N-succinimido, N-phthalimido, 2,5-dioxo-1-oxazolidinyl, 3-dodecyl-2,5-dioxo-1-imidazolyl, and N-acetyl-N-dodecylamino,

- ethoxycarbonylamino, phenoxycarbonylamino, benzyloxycarbonylamino, hexadecyloxycarbonylamino, 2,4-di-*t*-butylphenoxycarbonylamino, phenylcarbonylamino, 2,5-(di-*t*-pentylphenyl)carbonylamino, *p*-dodecylphenylcarbonylamino, *p*-tolylcarbonylamino, N-methylureido, N,N-dimethylureido, N-methyl-N-dodecylureido, N-hexadecylureido, N,N-dioctadecylureido, N,N-dioctyl-N'-ethylureido, N-phenylureido, N,N-diphenylureido, N-phenyl-N-*p*-tolylureido, N-(*m*-hexadecylphenyl)ureido, N,N-(2,5-di-*t*-pentylphenyl)-N'-ethylureido, and *t*-butylcarbonamido; sulfonamido, such as methylsulfonamido, benzenesulfonamido, *p*-tolylsulfonamido, *p*-dodecylbenzenesulfonamido, N-methyltetradecylsulfonamido, N,N-dipropylsulfamoylamino, and hexadecylsulfonamido; sulfamoyl, such as N-methylsulfamoyl, N-ethylsulfamoyl, N,N-dipropylsulfamoyl, N-hexadecylsulfamoyl, N,N-dimethylsulfamoyl; N-[3-(dodecyloxy)propyl]sulfamoyl, N-[4-(2,4-di-*t*-pentylphenoxy)butyl]sulfamoyl, N-methyl-N-tetradecylsulfamoyl, and N-dodecylsulfamoyl; carbamoyl, such as N-methylcarbamoyl, N,N-dibutylcarbamoyl, N-octadecylcarbamoyl, N-[4-(2,4-di-*t*-pentylphenoxy)butyl]carbamoyl, N-methyl-N-tetradecylcarbamoyl, and N,N-dioctylcarbamoyl; acyl, such as acetyl, (2,4-di-*t*-amylphenoxy)acetyl, phenoxycarbonyl, *p*-dodecyloxyphenoxycarbonyl methoxycarbonyl, butoxycarbonyl, tetradecyloxycarbonyl, ethoxycarbonyl, benzyloxycarbonyl, 3-pentadecyloxycarbonyl, and dodecyloxycarbonyl; sulfonyl, such as methoxysulfonyl, octyloxysulfonyl, tetradecyloxysulfonyl, 2-ethylhexyloxysulfonyl, phenoxysulfonyl, 2,4-di-*t*-pentylphenoxysulfonyl, methylsulfonyl, octylsulfonyl, 2-ethylhexylsulfonyl, dodecylsulfonyl, hexadecylsulfonyl, phenylsulfonyl, 4-nonylphenylsulfonyl, and *p*-tolylsulfonyl; sulfonyloxy, such as dodecylsulfonyloxy, and hexadecylsulfonyloxy; sulfinyl, such as methylsulfinyl, octylsulfinyl, 2-ethylhexylsulfinyl, dodecylsulfinyl, hexadecylsulfinyl, phenylsulfinyl, 4-nonylphenylsulfinyl, and *p*-tolylsulfinyl; thio, such as ethylthio, octylthio, benzylthio, tetradecylthio, 2-(2,4-di-*t*-pentylphenoxy)ethylthio, phenylthio, 2-butoxy-5-*t*-octylphenylthio, and *p*-tolylthio; acyloxy, such as acetyloxy, benzoyloxy, octadecanoyloxy, *p*-dodecylamidobenzoyloxy, N-phenylcarbamoyloxy, N-ethylcarbamoyloxy, and

10021408-1212001

cyclohexylcarbonyloxy; amine, such as phenylanilino, 2-chloroanilino, diethylamine, dodecylamine; imino, such as 1 (N-phenylimido)ethyl, N-succinimido or 3-benzylhydantoinyl; phosphate, such as dimethylphosphate and ethylbutylphosphate; phosphite, such as diethyl and dihexylphosphite; a  
5 heterocyclic group, a heterocyclic oxy group or a heterocyclic thio group, each of which may be substituted and which contain a 3 to 7 membered heterocyclic ring composed of carbon atoms and at least one hetero atom selected from the group consisting of oxygen, nitrogen and sulfur, such as 2-furyl, 2-thienyl, 2-benzimidazolyloxy or 2-benzothiazolyl; quaternary ammonium, such as  
10 triethylammonium; and silyloxy, such as trimethylsilyloxy.

If desired, the substituents may themselves be further substituted one or more times with the described substituent groups. The particular substituents used may be selected by those skilled in the art to attain the desired photographic properties for a specific application and can include, for example,  
15 hydrophobic groups, solubilizing groups, blocking groups, and releasing or releasable groups. When a molecule may have two or more substituents, the substituents may be joined together to form a ring such as a fused ring unless otherwise provided. Generally, the above groups and substituents thereof may include those having up to 48 carbon atoms, typically 1 to 36 carbon atoms and  
20 usually less than 24 carbon atoms, but greater numbers are possible depending on the particular substituents selected.

The materials useful in the invention can be used in any of the ways and in any of the combinations known in the art. Typically, the invention materials are incorporated in a melt and coated as a layer described herein on a  
25 support to form part of a photographic element. When the term "associated" is employed, it signifies that a reactive compound is in or adjacent to a specified layer where, during processing, it is capable of reacting with other components.

To control the migration of various components, it may be desirable to include a high molecular weight hydrophobe or "ballast" group in  
30 coupler molecules. Representative ballast groups include substituted or unsubstituted alkyl or aryl groups containing 8 to 48 carbon atoms. Representative substituents on such groups include alkyl, aryl, alkoxy, aryloxy,

alkylthio, hydroxy, halogen, alkoxycarbonyl, aryloxycarbonyl, carboxy, acyl, acyloxy, amino, anilino, carbonamido, carbamoyl, alkylsulfonyl, arylsulfonyl, sulfonamido, and sulfamoyl groups wherein the substituents typically contain 1 to 42 carbon atoms. Such substituents can also be further substituted.

5           The photographic elements can be single color elements or multicolor elements. Multicolor elements contain image dye-forming units sensitive to each of the three primary regions of the spectrum. Each unit can comprise a single emulsion layer or multiple emulsion layers sensitive to a given region of the spectrum. The layers of the element, including the layers of the  
10 image-forming units, can be arranged in various orders as known in the art. In an alternative format, the emulsions sensitive to each of the three primary regions of the spectrum can be disposed as a single segmented layer.

          A typical multicolor photographic element comprises a support bearing a cyan dye image-forming unit comprised of at least one red-sensitive  
15 silver halide emulsion layer having associated therewith at least one cyan dye-forming coupler, a magenta dye image-forming unit comprising at least one green-sensitive silver halide emulsion layer having associated therewith at least one magenta dye-forming coupler, and a yellow dye image-forming unit comprising at least one blue-sensitive silver halide emulsion layer having associated therewith at  
20 least one yellow dye-forming coupler. The element can contain additional layers, such as filter layers, interlayers, overcoat layers, and subbing layers.

          If desired, the photographic element can be used in conjunction with an applied magnetic layer as described in Research Disclosure, November 1992, Item 34390 published by Kenneth Mason Publications, Ltd., Dudley Annex,  
25 12a North Street, Emsworth, Hampshire PO10 7DQ, ENGLAND, and as described in Hatsumi Kyoukai Koukai Gihou No. 94-6023, published March 15, 1994, available from the Japanese Patent Office. When it is desired to employ the inventive materials in a small format film, Research Disclosure, June 1994, Item 36230, provides suitable embodiments.

30           In the following discussion of suitable materials for use in the emulsions and elements of this invention, reference will be made to Research Disclosure, September 1996, Item 38957, available as described above, which is

referred to herein by the term "Research Disclosure". The Sections hereinafter referred to are Sections of the Research Disclosure.

Except as provided, the silver halide emulsion containing elements employed in this invention can be either negative-working or positive-working as indicated by the type of processing instructions (i.e. color negative, reversal, or direct positive processing) provided with the element. Suitable emulsions and their preparation as well as methods of chemical and spectral sensitization are described in Sections I through V. Various additives such as UV dyes, brighteners, antifoggants, stabilizers, light absorbing and scattering materials, and physical property modifying addenda such as hardeners, coating aids, plasticizers, lubricants and matting agents are described, for example, in Sections II and VI through VIII. Color materials are described in Sections X through XIII. Suitable methods for incorporating couplers and dyes, including dispersions in organic solvents, are described in Section X(E). Scan facilitating is described in Section XIV. Supports, exposure, development systems, and processing methods and agents are described in Sections XV to XX. The information contained in the September 1994 Research Disclosure, Item No. 36544 referenced above, is updated in the September 1996 Research Disclosure, Item No. 38957. Certain desirable photographic elements and processing steps, including those useful in conjunction with color reflective prints, are described in Research Disclosure, Item 37038, February 1995.

Coupling-off groups are well known in the art. Such groups can determine the chemical equivalency of a coupler, i.e., whether it is a 2-equivalent or a 4-equivalent coupler, or modify the reactivity of the coupler. Such groups can advantageously affect the layer in which the coupler is coated, or other layers in the photographic recording material, by performing, after release from the coupler, functions such as dye formation, dye hue adjustment, development acceleration or inhibition, bleach acceleration or inhibition, electron transfer facilitation, and color correction.

The presence of hydrogen at the coupling site provides a 4-equivalent coupler, and the presence of another coupling-off group usually provides a 2-equivalent coupler. Representative classes of such coupling-off

groups include, for example, chloro, alkoxy, aryloxy, hetero-oxy, sulfonyloxy, acyloxy, acyl, heterocyclyl, sulfonamido, mercaptotetrazole, benzothiazole, mercaptopropionic acid, phosphonyloxy, arylthio, and arylazo. These coupling-off groups are described in the art, for example, in U.S. Pat. Nos. 2,455,169, 3,227,551, 3,432,521, 3,476,563, 3,617,291, 3,880,661, 4,052,212 and 4,134,766; and in UK. Patents and published application Nos. 1,466,728, 1,531,927, 1,533,039, 2,006,755A and 2,017,704A.

Image dye-forming couplers may be included in the element such as couplers that form cyan dyes upon reaction with oxidized color developing

- agents which are described in such representative patents and publications as: "Farbkuppler-eine Literature Übersicht," published in Agfa Mitteilungen, Band III, pp. 156-175 (1961) as well as in U.S. Patent Nos. 2,367,531; 2,423,730; 2,474,293; 2,772,162; 2,895,826; 3,002,836; 3,034,892; 3,041,236; 4,333,999; 4,746,602; 4,753,871; 4,770,988; 4,775,616; 4,818,667; 4,818,672; 4,822,729; 4,839,267; 4,840,883; 4,849,328; 4,865,961; 4,873,183; 4,883,746; 4,900,656; 4,904,575; 4,916,051; 4,921,783; 4,923,791; 4,950,585; 4,971,898; 4,990,436; 4,996,139; 5,008,180; 5,015,565; 5,011,765; 5,011,766; 5,017,467; 5,045,442; 5,051,347; 5,061,613; 5,071,737; 5,075,207; 5,091,297; 5,094,938; 5,104,783; 5,178,993; 5,813,729; 5,187,057; 5,192,651; 5,200,305; 5,202,224; 5,206,130; 5,208,141; 5,210,011; 5,215,871; 5,223,386; 5,227,287; 5,256,526; 5,258,270; 5,272,051; 5,306,610; 5,326,682; 5,366,856; 5,378,596; 5,380,638; 5,382,502; 5,384,236; 5,397,691; 5,415,990; 5,434,034; 5,441,863; EPO 0 246 616; EPO 0 250 201; EPO 0 271 323; EPO 0 295 632; EPO 0 307 927; EPO 0 333 185; EPO 0 378 898; EPO 0 389 817; EPO 0 487 111; EPO 0 488 248; EPO 0 539 034; EPO 0 545 300; EPO 0 556 700; EPO 0 556 777; EPO 0 556 858; EPO 0 569 979; EPO 0 608 133; EPO 0 636 936; EPO 0 651 286; EPO 0 690 344; German OLS 4,026,903; German OLS 3,624,777. and German OLS 3,823,049. Typically such couplers are phenols, naphthols, or pyrazoloazoles.

- Couplers that form magenta dyes upon reaction with oxidized color developing agent are described in such representative patents and publications as: "Farbkuppler-eine Literature Übersicht," published in Agfa Mitteilungen, Band III, pp. 126-156 (1961) as well as U.S. Patents 2,311,082 and 2,369,489;



- 2,343,701; 2,600,788; 2,908,573; 3,062,653; 3,152,896; 3,519,429; 3,758,309;  
3,935,015; 4,540,654; 4,745,052; 4,762,775; 4,791,052; 4,812,576; 4,835,094;  
4,840,877; 4,845,022; 4,853,319; 4,868,099; 4,865,960; 4,871,652; 4,876,182;  
4,892,805; 4,900,657; 4,910,124; 4,914,013; 4,921,968; 4,929,540; 4,933,465;  
5 4,942,116; 4,942,117; 4,942,118; U.S. Patent 4,959,480; 4,968,594; 4,988,614;  
4,992,361; 5,002,864; 5,021,325; 5,066,575; 5,068,171; 5,071,739; 5,100,772;  
5,110,942; 5,116,990; 5,118,812; 5,134,059; 5,155,016; 5,183,728; 5,234,805;  
5,235,058; 5,250,400; 5,254,446; 5,262,292; 5,300,407; 5,302,496; 5,336,593;  
5,350,667; 5,395,968; 5,354,826; 5,358,829; 5,368,998; 5,378,587; 5,409,808;  
10 5,411,841; 5,418,123; 5,424,179; EPO 0 257 854; EPO 0 284 240; EPO  
0 341 204; EPO 347,235; EPO 365,252; EPO 0 422 595; EPO 0 428 899; EPO  
0 428 902; EPO 0 459 331; EPO 0 467 327; EPO 0 476 949; EPO 0 487 081;  
EPO 0 489 333; EPO 0 512 304; EPO 0 515 128; EPO 0 534 703; EPO 0 554 778;  
EPO 0 558 145; EPO 0 571 959; EPO 0 583 832; EPO 0 583 834; EPO 0 584 793;  
15 EPO 0 602 748; EPO 0 602 749; EPO 0 605 918; EPO 0 622 672; EPO 0 622 673;  
EPO 0 629 912; EPO 0 646 841; EPO 0 656 561; EPO 0 660 177; EPO 0 686 872;  
WO 90/10253; WO 92/09010; WO 92/10788; WO 92/12464; WO 93/01523; WO  
93/02392; WO 93/02393; WO 93/07534; UK Application 2,244,053; Japanese  
Application 03192-350; German OLS 3,624,103; German OLS 3,912,265; and  
20 German OLS 40 08 067. Typically such couplers are pyrazolones,  
pyrazoloazoles, or pyrazolobenzimidazoles that form magenta dyes upon reaction  
with oxidized color developing agents.

- Couplers that form yellow dyes upon reaction with oxidized color  
developing agent are described in such representative patents and publications as:
- 25 "Farbkuppler-eine Literature Übersicht," published in Agfa Mitteilungen; Band  
III; pp. 112-126 (1961); as well as U.S. Patent 2,298,443; 2,407,210; 2,875,057;  
3,048,194; 3,265,506; 3,447,928; 4,022,620; 4,443,536; 4,758,501; 4,791,050;  
4,824,771; 4,824,773; 4,855,222; 4,978,605; 4,992,360; 4,994,361; 5,021,333;  
5,053,325; 5,066,574; 5,066,576; 5,100,773; 5,118,599; 5,143,823; 5,187,055;  
30 5,190,848; 5,213,958; 5,215,877; 5,215,878; 5,217,857; 5,219,716; 5,238,803;  
5,283,166; 5,294,531; 5,306,609; 5,328,818; 5,336,591; 5,338,654; 5,358,835;  
5,358,838; 5,360,713; 5,362,617; 5,382,506; 5,389,504; 5,399,474; 5,405,737;

5,411,848; 5,427,898; EPO 0 327 976; EPO 0 296 793; EPO 0 365 282; EPO 0 379 309; EPO 0 415 375; EPO 0 437 818; EPO 0 447 969; EPO 0 542 463; EPO 0 568 037; EPO 0 568 196; EPO 0 568 777; EPO 0 570 006; EPO 0 573 761; EPO 0 608 956; EPO 0 608 957; and EPO 0 628 865. Such couplers are typically

5 open chain ketomethylene compounds.

Couplers that form colorless products upon reaction with oxidized color developing agent are described in such representative patents as:

UK. 861,138; U.S. Pat. Nos. 3,632,345; 3,928,041; 3,958,993 and 3,961,959.

Typically such couplers are cyclic carbonyl containing compounds that form  
10 colorless products on reaction with an oxidized color developing agent.

Couplers that form black dyes upon reaction with oxidized color developing agent are described in such representative patents as U.S. Patent Nos. 1,939,231; 2,181,944; 2,333,106; and 4,126,461; German OLS No. 2,644,194 and German OLS No. 2,650,764. Typically, such couplers are resorcinols or *m*-  
15 aminophenols that form black or neutral products on reaction with oxidized color developing agent.

In addition to the foregoing, so-called "universal" or "washout" couplers may be employed. These couplers do not contribute to image dye-formation. Thus, for example, a naphthol having an unsubstituted carbamoyl or  
20 one substituted with a low molecular weight substituent at the 2- or 3- position may be employed. Couplers of this type are described, for example, in U.S. Patent Nos. 5,026,628, 5,151,343, and 5,234,800.

It may be useful to use a combination of couplers any of which may contain known ballasts or coupling-off groups such as those described in U.S.  
25 Patent 4,301,235; U.S. Patent 4,853,319 and U.S. Patent 4,351,897. The coupler may contain solubilizing groups such as described in U.S. Patent 4,482,629. The coupler may also be used in association with "wrong" colored couplers (e.g. to adjust levels of interlayer correction) and, in color negative applications, with masking couplers such as those described in EP 213.490; Japanese Published  
30 Application 58-172,647; U.S. Patent Nos. 2,983,608; 4,070,191; and 4,273,861; German Applications DE 2,706,117 and DE 2,643,965; UK. Patent 1,530,272;

and Japanese Application 58-113935. The masking couplers may be shifted or blocked, if desired.

Typically, couplers are incorporated in a silver halide emulsion layer in a mole ratio to silver of 0.05 to 1.0 and generally 0.1 to 0.5. Usually the  
5 couplers are dispersed in a high-boiling organic solvent in a weight ratio of solvent to coupler of 0.1 to 10.0 and typically 0.1 to 2.0 although dispersions using no permanent coupler solvent are sometimes employed.

The invention may be used in association with materials that release Photographically Useful Groups (PUGS) that accelerate or otherwise  
10 modify the processing steps e.g. of bleaching or fixing to improve the quality of the image. Bleach accelerator releasing couplers such as those described in EP 193,389; EP 301,477; U.S. 4,163,669; U.S. 4,865,956; and U.S. 4,923,784, may be useful. Also contemplated is use in association with nucleating agents, development accelerators or their precursors (UK Patent 2,097,140; UK. Patent  
15 2,131,188); electron transfer agents (U.S. 4,859,578; U.S. 4,912,025); antifogging and anti color-mixing agents such as derivatives of hydroquinones, aminophenols, amines, gallic acid; catechol; ascorbic acid; hydrazides; sulfonamidophenols; and non color-forming couplers.



The invention may also be used in combination with filter dye  
20 layers comprising colloidal silver sol or yellow, cyan, and/or magenta filter dyes, either as oil-in-water dispersions, latex dispersions or as solid particle dispersions. Additionally, they may be used with "smearing" couplers (e.g. as described in U.S. 4,366,237; EP 96,570; U.S. 4,420,556; and U.S. 4,543,323.) Also, the materials useful in the invention may be blocked or coated in protected form as  
25 described, for example, in Japanese Application 61/258,249 or U.S. 5,019,492.



The invention may further be used in combination with image-modifying compounds that release PUGS such as "Developer Inhibitor-Releasing" compounds (DIR's). DIR's useful in conjunction with the invention are known in the art and examples are described in U.S. Patent Nos. 3,137,578; 3,148,022;  
30 3,148,062; 3,227,554; 3,384,657; 3,379,529; 3,615,506; 3,617,291; 3,620,746; 3,701,783; 3,733,201; 4,049,455; 4,095,984; 4,126,459; 4,149,886; 4,150,228; 4,211,562; 4,248,962; 4,259,437; 4,362,878; 4,409,323; 4,477,563; 4,782,012;

4,962,018; 4,500,634; 4,579,816; 4,607,004; 4,618,571; 4,678,739; 4,746,600;  
4,746,601; 4,791,049; 4,857,447; 4,865,959; 4,880,342; 4,886,736; 4,937,179;  
4,946,767; 4,948,716; 4,952,485; 4,956,269; 4,959,299; 4,966,835; 4,985,336 as  
well as in patent publications GB 1,560,240; GB 2,007,662; GB 2,032,914; GB  
5 2,099,167; DE 2,842,063; DE 2,937,127; DE 3,636,824; DE 3,644,416 as well as  
the following European Patent Publications: 272,573; 335,319; 336,411; 346,  
899; 362, 870; 365,252; 365,346; 373,382; 376,212; 377,463; 378,236; 384,670;  
396,486; 401,612; 401,613.

Such compounds are also disclosed in "Developer-Inhibitor-  
10 Releasing (DIR) Couplers for Color Photography," C.R. Barr, J.R. Thirtle and  
P.W. Vittum in Photographic Science and Engineering, Vol. 13, p. 174 (1969).  
Generally, the developer inhibitor-releasing (DIR) couplers include a coupler  
moiety and an inhibitor coupling-off moiety (IN). The inhibitor-releasing  
couplers may be of the time-delayed type (DIAR couplers) which also include a  
15 timing moiety or chemical switch which produces a delayed release of inhibitor.  
Examples of typical inhibitor moieties are: oxazoles, thiazoles, diazoles, triazoles,  
oxadiazoles, thiadiazoles, oxathiazoles, thiatrazoles, benzotriazoles, tetrazoles,  
benzimidazoles, indazoles, isoindazoles, mercaptotetrazoles, selenotetrazoles,  
mercaptobenzothiazoles, selenobenzothiazoles, mercaptobenzoxazoles,  
20 selenobenzoxazoles, mercaptobenzimidazoles, selenobenzimidazoles,  
benzodiazoles, mercaptooxazoles, mercaptothiadiazoles, mercaptothiazoles,  
mercaptotriazoles, mercaptooxadiazoles, mercaptodiazoles, mercaptooxathiazoles,  
telteurotetrazoles or benzisodiazoles. In a preferred embodiment, the inhibitor  
moiety or group is selected from the following formulas:

25







wherein  $R_I$  is selected from the group consisting of straight and branched alkyls of from 1 to about 8 carbon atoms, benzyl, phenyl, and alkoxy groups and such groups containing none, one or more than one such substituent;  $R_{II}$  is selected from  $R_I$  and  $-SR_I$ ;  $R_{III}$  is a straight or branched alkyl group of from 1 to about 5 carbon atoms and  $m$  is from 1 to 3; and  $R_{IV}$  is selected from the group consisting of hydrogen, halogens and alkoxy, phenyl and carbonamido groups,  $-COOR_V$  and  $-NHCOR_V$  wherein  $R_V$  is selected from substituted and unsubstituted alkyl and aryl groups.

Although it is typical that the coupler moiety included in the developer inhibitor-releasing coupler forms an image dye corresponding to the layer in which it is located, it may also form a different color as one associated with a different film layer. It may also be useful that the coupler moiety included in the developer inhibitor-releasing coupler forms colorless products and/or products that wash out of the photographic material during processing (so-called "universal" couplers).



A compound such as a coupler may release a PUG directly upon reaction of the compound during processing, or indirectly through a timing or linking group. A timing group produces the time-delayed release of the PUG such groups using an intramolecular nucleophilic substitution reaction (U.S. 4,248,962); groups utilizing an electron transfer reaction along a conjugated system (U.S. 4,409,323; 4,421,845; 4,861,701, Japanese Applications 57-188035; 58-98728; 58-209736; 58-209738); groups that function as a coupler or reducing agent after the coupler reaction (U.S. 4,438,193; U.S. 4,618,571) and groups that combine the features describe above. It is typical that the timing group is of one of the formulas:













- wherein IN is the inhibitor moiety, R<sub>VII</sub> is selected from the group consisting of nitro, cyano, alkylsulfonyl, sulfamoyl; and sulfonamido groups; a is 0 or 1; and R<sub>VI</sub> is selected from the group consisting of substituted and unsubstituted alkyl and phenyl groups. The oxygen atom of each timing group is bonded to the coupling-off position of the respective coupler moiety of the DIAR.

- The timing or linking groups may also function by electron transfer down an unconjugated chain. Linking groups are known in the art under various names. Often they have been referred to as groups capable of utilizing a hemiacetal or iminoketal cleavage reaction or as groups capable of utilizing a cleavage reaction due to ester hydrolysis such as U.S. 4,546,073. This electron transfer down an unconjugated chain typically results in a relatively fast decomposition and the production of carbon dioxide, formaldehyde, or other low molecular weight by-products. The groups are exemplified in EP 523,451, U.S. 4,146,396, Japanese Kokai 60-249148 and 60-249149.

Suitable developer inhibitor-releasing couplers for use in the present invention include, but are not limited to, the following:







- It is also contemplated that the present invention may be employed to obtain reflection color prints as described in Research Disclosure, November 1979, Item 18716, available from Kenneth Mason Publications, Ltd, Dudley Annex, 12a North Street, Emsworth, Hampshire P0101 7DQ, England. Materials useful in the invention may be coated on pH adjusted support as described in U.S. 4,917,994; on a support with reduced oxygen permeability (EP 553,339); with epoxy solvents (EP 164,961); with nickel complex stabilizers (U.S. 4,346,165; U.S. 4,540,653 and U.S. 4,906,559 for example); with ballasted chelating agents such as those in U.S. 4,994,359 to reduce sensitivity to polyvalent cations such as calcium; and with stain reducing compounds such as described in U.S. 5,068,171. Other compounds useful in combination with the invention are disclosed in Japanese Published Applications described in Derwent Abstracts having accession



numbers as follows: 90-072,629, 90-072,630; 90-072,631; 90-072,632; 90-072,633; 90-072,634; 90-077,822; 90-078,229; 90-078,230; 90-079,336; 90-079,337; 90-079,338; 90-079,690; 90-079,691; 90-080,487; 90-080,488; 90-080,489; 90-080,490; 90-080,491; 90-080,492; 90-080,494; 90-085,928; 90-086,669; 90-086,670; 90-087,360; 90-087,361; 90-087,362; 90-087,363; 90-087,364; 90-088,097; 90-093,662; 90-093,663; 90-093,664; 90-093,665; 90-093,666; 90-093,668; 90-094,055; 90-094,056; 90-103,409; 83-62,586; 83-09,959.

Conventional radiation-sensitive silver halide emulsions can be employed in the practice of this invention. Such emulsions are illustrated by *Research Disclosure*, Item 38755, September 1996, I. Emulsion grains and their preparation.

Especially useful in this invention are tabular grain silver halide emulsions. Tabular grains are those having two parallel major crystal faces and having an aspect ratio of at least 2. The term "aspect ratio" is the ratio of the equivalent circular diameter (ECD) of a grain major face divided by its thickness (t). Tabular grain emulsions are those in which the tabular grains account for at least 50 percent (preferably at least 70 percent and optimally at least 90 percent) of the total grain projected area. Preferred tabular grain emulsions are those in which the average thickness of the tabular grains is less than 0.3 micrometer (preferably thin--that is, less than 0.2 micrometer and most preferably ultrathin--that is, less than 0.07 micrometer). The major faces of the tabular grains can lie in either {111} or {100} crystal planes. The mean ECD of tabular grain emulsions rarely exceeds 10 micrometers and more typically is less than 5 micrometers.

In their most widely used form tabular grain emulsions are high bromide {111} tabular grain emulsions. Such emulsions are illustrated by Kofron et al U.S. Patent 4,439,520, Wilgus et al U.S. Patent 4,434,226, Solberg et al U.S. Patent 4,433,048, Maskasky U.S. Patents 4,435,501,, 4,463,087 and 4,173,320, Daubendiek et al U.S. Patents 4,414,310 and 4,914,014, Sowinski et al U.S. Patent 4,656,122, Piggitt et al U.S. Patents 5,061,616 and 5,061,609, Tsaur et al U.S. Patents 5,147,771, '772, '773, 5,171,659 and 5,252,453, Black et al 5,219,720 and 5,334,495, Delton U.S. Patents 5,310,644, 5,372,927 and 5,460,934, Wen U.S.

Patent 5,470,698, Fenton et al U.S. Patent 5,476,760, Eshelman et al U.S. Patents 5,612,175 and 5,614,359, and Irving et al U.S. Patent 5,667,954.

5 Ultrathin high bromide {111} tabular grain emulsions are illustrated by Daubendiek et al U.S. Patents 4,672,027, 4,693,964, 5,494,789, 5,503,971 and 5,576,168, Antoniadis et al U.S. Patent 5,250,403, Olm et al U.S. Patent 5,503,970, Deaton et al U.S. Patent 5,582,965, and Maskasky U.S. Patent 5,667,955.

High bromide {100} tabular grain emulsions are illustrated by Mignot U.S. Patents 4,386,156 and 5,386,156.

10 High chloride {111} tabular grain emulsions are illustrated by Wey U.S. Patent 4,399,215, Wey et al U.S. Patent 4,414,306, Maskasky U.S. Patents 4,400,463, 4,713,323, 5,061,617, 5,178,997, 5,183,732, 5,185,239, 5,399,478 and 5,411,852, and Maskasky et al U.S. Patents 5,176,992 and 5,178,998. Ultrathin high chloride {111} tabular grain emulsions are illustrated by Maskasky U.S. Patents 5,271,858 and 5,389,509.

15 High chloride {100} tabular grain emulsions are illustrated by Maskasky U.S. Patents 5,264,337, 5,292,632, 5,275,930 and 5,399,477, House et al U.S. Patent 5,320,938, Brust et al U.S. Patent 5,314,798, Szajewski et al U.S. Patent 5,356,764, Chang et al U.S. Patents 5,413,904 and 5,663,041, Oyamada U.S. Patent 5,593,821, Yamashita et al U.S. Patents 5,641,620 and 5,652,088, Saitou et al U.S. Patent 5,652,089, and Oyamada et al U.S. Patent 5,665,530. Ultrathin high chloride {100} tabular grain emulsions can be prepared by nucleation in the presence of iodide, following the teaching of House et al and Chang et al, cited above.

25 The emulsions can be surface-sensitive emulsions, i.e., emulsions that form latent images primarily on the surfaces of the silver halide grains, or the emulsions can form internal latent images predominantly in the interior of the silver halide grains. The emulsions can be negative-working emulsions, such as surface-sensitive emulsions or unfogged internal latent image-forming emulsions, or direct-positive emulsions of the unfogged, internal latent image-forming type, which are positive-working when development is conducted with uniform light

30

exposure or in the presence of a nucleating agent. Tabular grain emulsions of the latter type are illustrated by Evans et al. U.S. 4,504,570.

Photographic elements can be exposed to actinic radiation, typically in the visible region of the spectrum, to form a latent image and can then  
5 be processed to form a visible dye image. Processing to form a visible dye image includes the step of contacting the element with a color-developing agent to reduce developable silver halide and oxidize the color-developing agent. Oxidized color developing agent in turn reacts with the coupler to yield a dye. If desired "Redox Amplification" as described in Research Disclosure XVIII(B)(5)  
10 may be used.

A "color negative element" utilizes negative-working silver halide and provides a negative image upon processing. A first type of such element is a capture element, which is a color negative film that is designed for capturing an image in negative form rather than for viewing an image. A second type of such  
15 an element is a direct-view element that is designed, at least in part, for providing a positive image viewable by humans.

In the capture element, speed (the sensitivity of the element to low light conditions) is usually critical to obtaining sufficient image in such elements. Such elements are typically silver bromoiodide emulsions coated on a transparent  
20 support and are sold packaged with instructions to process in known color negative processes such as the Kodak C-41 process as described in The British Journal of Photography Annual of 1988, pages 191-198. If a color negative film element is to be subsequently employed to generate a viewable projection print as for a motion picture, a process such as the Kodak BCN-2 process described in the  
25 H-24 Manual available from Eastman Kodak Co. may be employed to provide the color negative image on a transparent support. Color negative development times are typically 3' 15" or less and desirably 90 or even 60 seconds or less.

A direct-view photographic element is one which yields a color image that is designed for human viewing (1) by reflected light, such as a  
30 photographic paper print, (2) by transmitted light, such as a display transparency, or (3) by projection, such as a color slide or a motion picture print. These direct-view elements may be exposed and processed in a variety of ways. For example,

- paper prints, display transparencies, and motion picture prints are typically produced by digitally printing or by optically printing an image from a color negative onto the direct-viewing element and processing through an appropriate negative-working photographic process to give a positive color image. The
- 5 element may be sold packaged with instructions for digital printing or for processing using a color negative optical printing process, for example the Kodak RA-4 process, as generally described in PCT WO 87/04534 or U.S. 4,975,357, to form a positive image. Color projection prints may be processed, for example, in accordance with the Kodak ECP-2 process as described in the H-24 Manual.
- 10 Color print development times are typically 90 seconds or less and desirably 45 or even 30 seconds or less. Color slides may be produced in a similar manner but are more typically produced by exposing the film directly in a camera and processing through a reversal color process or a direct positive process to give a positive color image. The foregoing images may also be produced by alternative processes
- 15 such as digital printing.

- Each of these types of photographic elements has its own particular requirements for dye hue, but in general they all require cyan dyes whose absorption bands are less deeply absorbing (that is, shifted away from the red end of the spectrum) than color negative films. This is because dyes in direct-view
- 20 elements are selected to have the best appearance when viewed by human eyes, whereas the dyes in image capture materials are designed to best match the needs of the printing process.

- A reversal element is capable of forming a positive image without optical printing. To provide a positive (or reversal) image, the color development
- 25 step is preceded by development with a non-chromogenic developing agent to develop exposed silver halide, but not form dye, and followed by uniformly fogging the element to render unexposed silver halide developable. Such reversal elements are typically sold packaged with instructions to process using a color reversal process such as the Kodak E-6 process as described in The British Journal
- 30 of Photography Annual of 1988, page 194. Alternatively, a direct positive emulsion can be employed to obtain a positive image.

The above elements are typically sold with instructions to process using the appropriate method such as the mentioned color negative (Kodak C-41), color print (Kodak RA-4), or reversal (Kodak E-6) process.

The photographic element of the invention can be incorporated into exposure structures intended for repeated use or exposure structures intended for limited use, variously referred to by names such as "single use cameras", "lens with film", or "photosensitive material package units".

Preferred color developing agents are *p*-phenylenediamines such as:

- 4-amino-N,N-diethylaniline hydrochloride,
- 4-amino-3-methyl-N,N-diethylaniline hydrochloride,
- 4-amino-3-methyl-N-ethyl-N-(2-methanesulfonamidoethyl)aniline sesquisulfate hydrate,
- 4-amino-3-methyl-N-ethyl-N-(2-hydroxyethyl)aniline sulfate,
- 4-amino-3-(2-methanesulfonamidoethyl)-N,N-diethylaniline hydrochloride, and
- 4-amino-N-ethyl-N-(2-methoxyethyl)-*m*-toluidine di-*p*-toluene sulfonic acid.


Development is usually followed by the conventional steps of bleaching, fixing, or bleach-fixing, to remove silver or silver halide, washing, and drying.

### Synthesis

The synthesis of oligomeric N-substituted glycines has been described by R.N. Zuckermann, et. al., *J. Am. Chem. Soc.*, **1992**, 10646-10647; and *J. Med. Chem.* **1994**, 2678-2685, by Eric M. Gordon and James F. Kerwin, Jr., *Combinatorial Chemistry and Molecular Diversity in Drug Discovery*, Wiley-Liss Press, John Wiley and Sons, Inc. New York, NY, **1998**, by Barry A. Bunin, *The Combinatorial Index*, Academic Press, New York, **1998**, and by Anthony W. Czarnik and Sheila DeWitt, *A Practical Guide to Combinatorial Chemistry*, ACS Professional Reference Books, Washington, D.C., **1997**. These materials have

been named peptoids because they are structurally related to the well known peptides. Further, S.M. Miller, et. al., *Bioorganic & Medicinal Chemistry Lett.*, **1994**, 2657-2662, has reported that N-substituted glycines are less rapidly hydrolyzed than similar L-amino containing peptides. Further it is anticipated that the unwanted hydrogen bonding of peptide linkages resulting in excessive aqueous solubility and crystallinity would be greatly diminished by introduction of a substituent on the nitrogen of the glycine fragment resulting in overall higher hydrophobicity which is a desirable characteristic of oil soluble imaging dyes. Peptoid substituted pyrazoloazole couplers may be synthesized by using a substituted pyrazoloazole in which the substituent on a carbon center contains a primary amine functional group.

Couplers of the present invention can be prepared by simple reactions and methods described below. U.S. Patent No. 5,256,526 provides useful intermediates for pyrrolotriazoles. Typical syntheses are illustrated by the following examples. In the examples below the starting material Rink Amide Resin, RAR, is commercially available from Argonaut Technologies, Inc., San Carlos, California, U.S.A. and is sold as the Fmoc protected material under the name of "PS-Rink-Fmoc Resin". This resin is a polystyrene backbone lightly (1-2%) cross-linked with divinylbenzene that has been functionalized with 4-(2,4-dimethoxyphenyl-Fmoc-aminomethyl)-phenylacetic acid, where the Fmoc group is 9-fluorenylmethyloxycarbonyl. The typical bead size is 90-105 microns and the typical amine loading is 0.90 - 0.95 mmol per gram. Similar Rink Amide Resins may also be sold from the same supplier under the trade names of Argogel<sup>TM</sup> and Argopore<sup>TM</sup> and will have bead size and amine loading which is different than "PS-Rink-Fmoc Resin". All described reactions were run in an Argonaut Quest 210<sup>TM</sup> Reactor using 5 mL Teflon reaction vessels which allowed for the simultaneous synthesis of up to twenty materials. It will be understood by one skilled in the art of organic synthesis that the reactions could also be carried out using conventional laboratory glassware with the expectation that similar results would be obtained.




### General Procedure Rink-NH-Fmoc Resin to Rink-NH<sub>2</sub> Resin

Into a 5mL reaction vessel was placed 0.2 grams of PS-Rink-Fmoc Resin. The resin was swelled by the addition of 4 mL of dimethylformamide.

- 5 The resin was agitated for 10 minutes and then the reaction vessel drained. The 9-fluorenylmethyloxycarbonyl (Fmoc) group was removed by treatment with 4 mL of a stock solution of 20% piperidine in dimethylformamide (volume %), agitated for 5 minutes, then the vessel was drained. This treatment with 20% piperidine in dimethylformamide was repeated once more. The resin was then washed four
- 10 times each with 4 mL of dimethylformamide, dichloromethane, tetrahydrofuran and finally with dimethylformamide again with 3 minutes agitation during each wash. The resin was air dried, and a sample removed from the reaction vessel and analyzed for percent Fmoc removal by NMR which showed complete removal of the Fmoc group.

15

### Example 1: Synthesis of Coupler A1



Step 1. Into a 5 mL reaction vessel containing 0.2 grams of Rink-NH<sub>2</sub> resin, described above, was added 3 mL of 0.6 Molar solution of bromoacetic acid

in dimethylformamide and 1 mL of 3.2 Molar solution of diisopropylcarbodiimide in dimethylformamide. The vessel contents were mixed at room temperature for 30 minutes, drained and the operation repeated twice more. The resin was then washed four times each with 4 mL of dimethylformamide and 4 mL of dimethylsulfoxide with 2.5 minutes agitation during each wash. After draining the wash solvent, a 4 mL solution of 2.0 Molar cyclohexylamine in dimethylsulfoxide was added. The resin reaction mixture was agitated for 18 hours at room temperature then drained. The resin was then washed eight times with 4 mL dimethylformamide each for 2.5 minutes agitation.

Step 2. After draining the wash solvent, was added 3 mL of 0.6 Molar solution of bromoacetic acid in dimethylformamide and 1 mL of 3.2 Molar solution of diisopropylcarbodiimide in dimethylformamide. The vessel contents were mixed for 30 minutes, drained and the operation repeated twice more. The resin was then washed four times each with 4 mL of dimethylformamide and 4 mL of dimethylsulfoxide with 2.5 minutes agitation during each wash. After draining the wash solvent, was added 4 mL of a 0.5 Molar dichloromethane, or optionally dimethylsulfoxide, solution of 6-(1,1-dimethylethyl)- $\alpha,\alpha,\beta,\beta$ -tetramethyl-1H-pyrazolo[5,1-c]-1,2,4-triazole-3-ethanamine, Chemical Abstracts Registry Number 210405-85-9 whose preparation has been described in Harder, et. al., US 5,925,503 (7/1999). The resin reaction mixture was agitated for 18 hours at room temperature then drained. The resin was then thoroughly washed four times each with 4 mL of dimethylformamide, tetrahydrofuran, methanol and dichloromethane with 2.5 minutes agitation during each wash. This process was repeated twice.

Step 3. After draining the wash solvent, the product was cleaved from the resin by treatment with 4 mL of 20% trifluoroacetic acid in dichloromethane (volume %), agitation for 30 min, then draining with collection of the cleavage solution containing product. This operation was repeated twice more and the collected cleavage solutions were combined. The resin was then washed three times with 4 mL of dichloromethane and this solvent was also added to the cleavage solution containing product. The cleavage solution was then concentrated to dryness under vacuum without heat to afford 117 milligrams of



product **A1** with 84% purity as determined by Reverse Phase High Pressure Liquid Chromatography analysis. Electrospray MS:  $[MH]^+ = 460m/e$ .

Using the above procedure with substitution of 2-ethylhexylamine, dodecylamine, 3-ethoxypropylamine, cyclohexanemethylamine, benzylamine, aniline or 1-(3-aminopropyl)-2-pyrrolidinone, and hexylamine all available from the Aldrich Chemical Company, Milwaukee, Wisconsin, U.S.A., for cyclohexylamine the peptoid couplers **A2** – **A8** could be obtained. Product weight, product purity and Electrospray mass spectrum  $[MH]^+$  ion are shown in Table I for couplers A1-A3.

Using the above procedure with substitution of 6-(1,1-dimethylethyl)-pyrazolo[5,1-c]-1,2,4-triazole-3-ethanamine, whose preparation as the toluenesulfonic acid salt, Chemical Abstract Registry Number 152602-81-8, is described in Kirschke, et.al., *Liebigs Ann. Chem.* **1994**, *10*, 1037-42, for 6-(1,1-dimethylethyl)- $\alpha,\alpha,\beta,\beta$ -tetramethyl-1H-pyrazolo[5,1-c]-1,2,4-triazole-3-ethanamine, the peptoid couplers **A'1** – **A'8** could be obtained.

Using the above procedure with substitution of 6-(1,1-dimethylethyl)- $\beta$ -methyl-pyrazolo[1,5-b]-1,2,4-triazole-2-ethanamine, Chemical Abstract Registry Number 180483-99-2, whose preparation is described in Kawagishi, T., et. al., US 5,605,788, for 6-(1,1-dimethylethyl)- $\alpha,\alpha,\beta,\beta$ -tetramethyl-1H-pyrazolo[5,1-c]-1,2,4-triazole-3-ethanamine, , and hexylamine, the peptoid couplers **E1** – **E8** could be obtained.

Identification of the products by Electrospray mass spectrometry consistently showed formation of the desired materials. Purification and isolation of the resultant couplers was found to be most conveniently achieved by Preparative High Pressure Liquid Chromatography as described below. Using these techniques materials of high purity could be routinely obtained.

Preparative High Pressure Liquid Chromatography – Mass Spectrometry of peptoid coupler combinatorial libraries.

The instrumentation employed for the purification was a Waters-Micromass Preparative HPLC-MS consisting of the following components: a Waters 4000 Preparative HPLC pump, a Gilson 215 Liquid Handler as Autoinjector with 208 racks containing 50 mL Falcon tubes for injection, a second

Gilson 215 Liquid Handler as Fraction Collector, a Waters 515 Pump for make up liquid and diluent into a splitter for analytical sampling, a LC Packings custom 1:1000 Preparative Splitter-Diluter for 20-70 mL per min flow rate, and Upchurch adjustable 5:1 splitter, a Waters 996 Photo Diode Array Detector, and a


5 Micromass ZMD Mass Spectrometer with Z-spray design and electrospray probe. This instrumentation was controlled by Mass Lynx Software for methods and settings, Fraction Lynx Software for fraction collection, and Open Lynx software for reporting data.

- Samples of crude peptoid products from solid phase synthesis were
- 10 dissolved in 2 mL of methylene chloride and injected automatically into the flow stream. The column for preparative HPLC was a Waters 4.0 x 10 cm Deltabond, 15  $\mu$ m, radial compression column held in a RCM module. Separations were achieved by the following general gradient method, where A = pure H<sub>2</sub>O, B = buffer (0.1 % acetic acid), C = acetonitrile, D = isopropanol with the flow rate of
- 15 30 mL/minute for each step. Keeping B constant at 10%, C was ramped from 75% to 100% in the first 10 minutes, then the solvent strength was increased further by ramping to 50% C and 50% D in the next 10 minutes. This was followed by a recycling step as part of the method to enable immediate injection of the next sample. This or similar gradient methods. The flow rate was 30 mL/minute for
- 20 each step.
- Gradient methods were incorporated into a general combinatorial separation method (sample list in Mass Lynx) for the purification of combinatorial samples in sequential order. The gradient method was designed to facilitate maximum separation of components in the reaction mixtures, and included column wash and
- 25 equilibration steps to allow for rapid recycling to allow for automated injection of multiple samples. By this method a new compound could be purified every 30 minutes.

- Mass spectral and UV-vis data was accumulated during each preparative HPLC run, including during the particular time of fractionation for the
- 30 component of interest. This method ensured isolation of pure products only (generally over 95 %) for each sample in rapid sequential order. This data could be reported by either manually selecting data for that period of time during a

fraction or peak, or by automatically generated for each peak by Open Lynx software, which generated a report summarizing chromatographic and spectral data.

## 5 Example 2: Synthesis of Coupler B1



- Step 1. Into a 5 mL reaction vessel containing 0.2 grams of Rink-NH<sub>2</sub> resin, described above, was added 3 mL of 0.6 Molar solution of bromoacetic acid in dimethylformamide and 1 mL of 3.2 Molar solution of diisopropylcarbodiimide in dimethylformamide. The vessel contents were mixed at room temperature for 30 minutes, drained and the operation repeated twice more. The resin was then washed eight times with 4 mL of dimethylformamide with 3 minutes agitation during each wash. After draining the wash solvent, a 4 mL solution of 2.0 Molar cyclohexylamine in dimethylsulfoxide was added. The resin reaction mixture was agitated for 18 hours at room temperature then drained. The resin was then washed eight times with 4 mL dimethylformamide each for 3 minutes agitation.

- Step 2. After draining the wash solvent, was added 3 mL of 0.6 Molar solution of bromoacetic acid in dimethylformamide and 1 mL of 3.2 Molar solution of diisopropylcarbodiimide in dimethylformamide. The vessel contents were mixed at room temperature for 30 minutes, drained and the operation repeated twice more. The resin was then washed eight times each with 4 mL of dimethylformamide with 3 minutes agitation during each wash. After draining the

wash solvent, a 4 mL solution of 2.0 Molar cyclohexylamine in dimethylsulfoxide was added. The resin reaction mixture was agitated for 18 hours at room temperature then drained. The resin was then washed eight times with 4 mL dimethylformamide each for 3 minutes agitation.

- 5       Step 3. After draining the wash solvent, was added 3 mL of 0.6 Molar solution of bromoacetic acid in dimethylformamide and 1 mL of 3.2 Molar solution of diisopropylcarbodiimide in dimethylformamide. The vessel contents were mixed for 30 minutes, drained and the operation repeated twice more. The resin was then washed eight times each with 4 mL of dimethylformamide with 3
- 10     minutes agitation during each wash. After draining the wash solvent, was added 4 mL of a 0.5 Molar dichloromethane, or optionally dimethylsulfoxide, solution of 6-(1,1-dimethylethyl)- $\alpha,\alpha,\beta,\beta$ -tetramethyl-1H-pyrazolo[5,1-c]-1,2,4-triazole-3-ethanamine, Chemical Abstracts Registry Number 210405-85-9 whose preparation has been described in Harder, et. al., US 5,925,503 (7/1999). The
- 15     resin reaction mixture was agitated for 18 hours at room temperature then drained. The resin was then thoroughly washed eight times each with 4 mL of dimethylformamide, then eight times 4 mL of dichloromethane with 3 minutes agitation during each wash.

- 20     Step 4. After draining the wash solvent, the product was cleaved from the resin by treatment with 4 mL of 20% trifluoroacetic acid in dichloromethane (volume %), agitation for 30 min, then draining with collection of the cleavage solution containing product. This operation was repeated twice more and the collected cleavage solutions were combined. The resin was then washed twice with 4 mL of dichloromethane and this solvent was also added to the cleavage
- 25     solution containing product. The cleavage solution was then concentrated to dryness under vacuum without heat to afford 126 milligrams of product **B1** with 97% purity as determined by Reverse Phase High Pressure Liquid Chromatography analysis. Electrospray MS: [MH]<sup>+</sup> = 599m/e.

- 30     Using the above procedure with substitution of 2-ethylhexylamine, dodecylamine, 3-ethoxypropylamine, cyclohexanemethylamine, benzylamine, and hexylamine, all available from the Aldrich Chemical Company, Milwaukee, Wisconsin, U.S.A., for cyclohexylamine the peptoid couplers **B2 – B25** could be

obtained. Product weight, product purity and Electrospray mass spectrum [MH]<sup>+</sup> ion are shown in Table I for couplers B1-B25.

Using the above procedure with substitution of 6-(1,1-dimethylethyl)-pyrazolo[5,1-c]-1,2,4-triazole-3-ethanamine, whose preparation as the toluenesulfonic acid salt, Chemical Abstract Registry Number 152602-81-8, is described in Kirschke, et al., *Liebigs Ann. Chem.* **1994**, *10*, 1037-42, for 6-(1,1-dimethylethyl)- $\alpha,\alpha,\beta,\beta$ -tetramethyl-1H-pyrazolo[5,1-c]-1,2,4-triazole-3-ethanamine, the peptoid couplers **B\*1** – **B\*25** could be obtained.

It will be understood by one skilled in the art that repetition of Step 2 in the procedure above would result in a five step procedure for the preparation of Inventive Coupler **C1**. In this manner, substitution of 2-ethylhexylamine, dodecylamine, 3-ethoxypropylamine, cyclohexanemethylamine, benzylamine, and hexylamine, all available from the Aldrich Chemical Company, Milwaukee, Wisconsin, U.S.A., for cyclohexylamine the inventive couplers **C2** – **C25** could also be obtained.

Substitution of N-[2-[6-(1,1-dimethylethyl)-1H-pyrazolo[5,1-c]-1,2,4-triazol-3-yl]-1,1,2-trimethylpropyl]-2-[2-aminophenoxy]-acetamide, which may be prepared by methods described in Romanet, et. al., US 5,972,587 (10/99), for 6-(1,1-dimethylethyl)- $\alpha,\alpha,\beta,\beta$ -tetramethyl-1H-pyrazolo[5,1-c]-1,2,4-triazole-3-ethanamine, in the above procedure Step 3 will result in the preparation of Inventive Coupler **D17**. Further, in this manner, substitution of 2-ethylhexylamine, dodecylamine, 3-ethoxypropylamine, cyclohexanemethylamine, benzylamine, and hexylamine, all available from the Aldrich Chemical Company, Milwaukee, Wisconsin, U.S.A., for cyclohexylamine the inventive couplers **D1** – **D22** could also be obtained. Product weight, product purity and Electrospray mass spectrum [MH]<sup>+</sup> ion are shown in Table I for inventive couplers D1-D8.

Table I.

Inventive Coupler	Product weight	Product purity	ES/MS [MH] <sup>+</sup>
A1 <sup>a</sup>	117 mg	84%	460
A2 <sup>a</sup>	96 mg	93%	490
A3 <sup>a</sup>	102 mg	92%	546
B1 <sup>a</sup>	126 mg	97%	599
B2 <sup>a</sup>	147 mg	98%	630
B3 <sup>a</sup>	92 mg	93%	686
B4 <sup>a</sup>	116 mg	97%	630
B5 <sup>a</sup>	152 mg	97%	660
B6 <sup>a</sup>	112 mg	95%	716
B7 <sup>a</sup>	114 mg	95%	686
B8 <sup>a</sup>	87 mg	93%	716
B9 <sup>a</sup>	73 mg	93%	772
B10 <sup>b</sup>	24 mg	95%	604
B11 <sup>b</sup>	39 mg	98%	606
B12 <sup>b</sup>	53 mg	95%	616
B13 <sup>b</sup>	51 mg	91%	610
B14 <sup>b</sup>	38 mg	99%	606
B15 <sup>b</sup>	56 mg	95%	608
B16 <sup>b</sup>	52 mg	98%	618
B17 <sup>b</sup>	61 mg	97%	612
B18 <sup>b</sup>	73 mg	98%	616
B19 <sup>b</sup>	74 mg	97%	618
B20 <sup>b</sup>	67 mg	98%	628
B21 <sup>b</sup>	70 mg	97%	622
B22 <sup>b</sup>	63 mg	98%	610
B23 <sup>b</sup>	77 mg	97%	612
B24 <sup>b</sup>	107 mg	98%	622
B25 <sup>b</sup>	53 mg	98%	616

Inventive Coupler	Product weight	Product purity	ES/MS [MH] <sup>+</sup>
D1 <sup>b</sup>	85 mg	98%	753
D2 <sup>b</sup>	96 mg	98%	755
D3 <sup>b</sup>	125 mg	98%	765
D4 <sup>b</sup>	105 mg	98%	759
D5 <sup>b</sup>	100 mg	98%	755
D6 <sup>b</sup>	102 mg	97%	757
D7 <sup>b</sup>	106 mg	98%	767
D8 <sup>b</sup>	93 mg	97%	761

Note: a. Crude product weight and crude product purity reported

Note: b. Isolated product weight and isolated product purity reported

- 5 It will be recognized by one skilled in the arts of imaging science that dyes of the current invention may be used in such technologies as thermal dye transfer and ink jet imaging, but are in no wise limited to those technologies. Dyes of the present invention can be prepared by reactions and methods described below. Typical syntheses are illustrated by the following examples.

## 10 Method of Dye Preparation

There are two methods used to prepare the dyes:

1. A small amount of dry coupler is weighed into a tube with the addition of a dry developer. An organic solvent such as acetonitrile or other common reversed-phase chromatographic solvent is added and the solution sonicated until the coupler has dissolved. An aliquot of aqueous base is added to both dissolve the developer and ionize the coupler. An oxidizing agent such as sodium persulfate is added to form the dye in solution. The total volume of around 20ml is rapidly transferred to a reversed-phase low pressure chromatographic system where the dye is eventually eluted by applying a solvent gradient. The effluent is monitored using a spectrophotometric detector tuned to an appropriate wavelength. The dye peak is collected and the solvent evaporated. Recovery of dye is usually in the 80%+ efficiency with purities generally averaging better than 98%. The recovered dye is analyzed by mass spectrometry

for identity and High Performance Liquid Chromatography for purity prior to making an emulsion.

2. The coupler is used to make the emulsion and the emulsion is coated with additional gelatin and small amounts of silver halide dispersion as described in the dispersion and coating sections. The coupler coating is converted to dye proportional to the amount of oxidant (in this method exposed silver halide) using conventional photographic development. This involves adding photographic developer solution for 1-2 minutes. A water rinse is done after removal of the developer. An aliquot of photographic bleach solution is added for 1 minute to remove any silver halide and then removed. Finally, repeated water rinses remove all residual developing chemicals.

#### Method of Dye Dispersion Preparation

- Many pertinent photographic properties such as hue, bandwidth, and light stability are hugely affected by the chemical formulations of the emulsions which are then coated onto various substrates. The purified dye or coupler is formulated into an emulsion using basically the same process.

1. The initial dye or coupler is weighed and dissolved in an organic solvent for easier transfer. An aliquot is added to a 15mL polypropylene centrifuge tube with the appropriate coupler solvent. Any formulation addenda such as stabilizers are also added at this time. The organic solvent is evaporated leaving the dye or coupler dissolved in a small pellet of coupler solvent. An additional auxiliary coupler solvent may be added with sonication or heating depending on solubility characteristics of the dye or coupler. The final components consisting of water, surfactant, and gelatin are added to make somewhere on the order of 4 ml of solution. This solution is sonicated using a 1/4inch probe with 150 watts of power at 20KHz for up to 25 minutes. The resulting emulsion has been shown to have median particle sizes on the order of 0.3 microns. This emulsion is filtered using 0.45 um filters. An aliquot is diluted with water and measured spectrophotometrically to verify the quality. The resulting formulation contains a gelatin concentration between 1-3.5%, dye



concentrations of 2-3 millimolar or coupler concentrations 6-14 millimolar. These emulsions are chilled and stored prior to coating

#### Method of Dye Dispersion Ink Jet Coating

- 5                   There are three distinct coating techniques which are used for micro-scale evaluations. Each of these techniques has its benefits and drawbacks. All of the techniques use conventional disposable microtiterplates or special holders which conform to titerplate dimensions. This allows us to use commercially available equipment for both measurement and photographic  
10 development.

1.       The first technique developed uses a special holder called a fadeplate. This holder accepts various web materials for laying down emulsions using ink-jet or similar technology. Typically a gelatin subbed Estar™ material is held in the fadeplate. The material is held down on a chilled vacuum plate when  
15 depositing material. Various devices are used to create patterns of dye emulsion dots which are merged by repeatedly dispensing a pattern with a slight offset. Typically dots are formed from drops of 100 pl to 50nl. The resulting patterned laydown is allowed to dry at room temperature before measurement of photographic parameters.

- 20       2.       The second technique involves adding 1-3% gelatin with sufficient dye emulsion and hardener to form 40 -60 ul of solution. After mixing, the plate is chill set and then allowed to dry overnight. The resulting plate has a uniformly coated mid section adequate for measurement.

3.       The third technique uses a mixture of coupler dispersion, gelatin,  
25 hardener, and silver dispersion. The silver dispersion is metered in using ink-jet technology and the plate mixed.. After mixing, the plate is chill set and then allowed to dry overnight. The plate is processed as described in the dye preparation section to form a uniform layer of dye for studies. This technique provides a coating of dye in a matrix which is very close to that found in  
30 conventional films and papers.


# Method of Coated Dye Light Fade Measurement

There are two major ways which are used to measure Light stability.


1. The fadeplates or titerplates are read spectrophotometrically using a commercial instrument. The spectra are stored in a database with specific information as to composition and exposure for each distinct section of the plate. The plates are mounted on white reflective cards, and a UV filter sheet is taped into place to screen the UV wavelengths. These cards are mounted in a large temperature controlled sphere with a 50 Klux light source revolving in the center. The plates are removed periodically and re-measured until the measurements indicate that about 10% of the dye has been lost. The data is then used to calculate light fade stability metrics. This typically takes on the order of 3 weeks but several plates can be faded simultaneously.

2. A more rapid method can be used with fadeplate type coatings. The coated section is cut out using a razor knife and placed in a modified holder resembling a standard cuvette. A specially modified spectrophotometer has been programmed to expose several cells at a time with 200Klux of light and periodically collect visible spectra. The data can be plotted and photographic parameters measured in a few hours to days.

The absorbance maximum of the coated dyes of this invention and the relative light fade of the coated dyes of this invention are shown in Table I and are compared to the corresponding data for Comparison Dyes **CD-1** and **CD-2** whose structures are shown below.



Comparison Dye **CD-1**



Comparison Dye **CD-2**

Table II. The invention dyes G1-G12 were formulated as dye dispersions with a stabilizer set. The invention dyes G13 to G28 were formulated as coupler dispersions with stabilizer sets.

Dye	<u>Absorbance</u> <u>maximum</u> Coated Image	<u>Half Band</u> <u>width</u> Coated Image	Relative light fade
Comparison Dye CD-1	545nm	90nm	2.78
Comparison Dye CD-2	549 nm	74 nm	1.00
Invention Dye G1	550 nm	84 nm	4.09
Invention Dye G2	550 nm	76 nm	2.71
Invention Dye G3	550nm	77 nm	2.51
Invention Dye G4	550 nm	79 nm	2.64
Invention Dye G5	550 nm	78 nm	1.99
Invention Dye G6	550 nm	80 nm	2.06
Invention Dye G7	550 nm	79 nm	3.53
Invention Dye G8	550 nm	78 nm	1.87
Invention Dye G9	550 nm	78 nm	2.41
Invention Dye G10	550 nm	79 nm	2.32
Invention Dye G11	550 nm	77 nm	2.13
Invention Dye G12	550 nm	78 nm	1.84
Invention Dye G13	555 nm	84 nm	1.80
Invention Dye G14	556 nm	77 nm	1.48
Invention Dye G15	554 nm	78 nm	1.39
Invention Dye G16	554 nm	78 nm	1.09
Invention Dye G17	555 nm	76 nm	1.23
Invention Dye G18	556 nm	76 nm	1.22
Invention Dye G19	555 nm	76 nm	1.13
Invention Dye G20	554 nm	75 nm	2.01
Invention Dye G21	556 nm	77 nm	1.15
Invention Dye G22	556 nm	76 nm	0.92
Invention Dye G23	555 nm	76 nm	1.08

Dye	Absorbance maximum Coated Image	Half Band width Coated Image	Relative light fade
Invention Dye G24	555 nm	76 nm	0.96
Invention Dye G25	554 nm	76 nm	0.94
Invention Dye G26	556 nm	74 nm	0.95
Invention Dye G27	556 nm	76 nm	1.04
Invention Dye G28	555 nm	76 nm	0.94

The data in Table II indicate that the dyes prepared from the couplers according to the present invention offer good light stability with a deeper absorbance maximum and without unwanted excessive absorbance band shape broadening compared with dye from pyrazolotriazole couplers of the prior art.

The invention has been described in detail with particular reference to certain preferred embodiments thereof, but it will be understood that variations and modifications can be effected within the spirit and scope of the invention.

The entire contents of the patents and other publications referred to in this specification are incorporated herein by reference.